








INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 215/54, A61K 31/63, 31/47, 31/44, C07D 401/12, 471/04, 513/04, 498/04, 495/04, C07F 9/547, A61K 31/66 // (C07D 513/04, 277:00, 221:00) (C07D 498/04, 261:00, 221:00) (C07D 471/04, 231:00, 221:00) (C07D 513/04, 275:00, 221:00) (C07D 471/04, 221:00, 221:00) (C07D 471/04, 235:00, 221:00)</p>	A1	<p>(11) International Publication Number: WO 99/18076</p>
<p>(21) International Application Number: PCT/US98/07380</p>		<p>(74) Agents: ALICE, Ronald, W.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.</p>
<p>(22) International Filing Date: 14 April 1998 (14.04.98)</p>		<p>(43) International Publication Date: 15 April 1999 (15.04.99)</p>
<p>(30) Priority Data: 08/944,188 6 October 1997 (06.10.97) US 09/055,856 6 April 1998 (06.04.98) US</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p>
<p>(71) Applicant: AMERICAN CYANAMID COMPANY [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).</p>		<p>Published With international search report.</p>
<p>(72) Inventors: LEVIN, Jeremy, Ian; 190 Treetop Circle, Nanuet, NY 10954 (US). ZASK, Arie; 200 Central Park South, New York, NY 10019 (US). GU, Yansong; 82 North Magnolia Avenue, Pearl River, NY 10965 (US). ALBRIGHT, Jay, Donald; 5 Clifford Court, Nanuet, NY 10954 (US). DU, Xuemei; 130 Sierra Vista Lane, Valley Cottage, NY 10989 (US).</p>		

(54) Title: THE PREPARATION AND USE OF ORTHO-SULFONAMIDO BICYCLIC HETEROARYL HYDROXAMIC ACIDS AS MATRIX METALLOPROTEINASE AND TACE INHIBITORS

(57) Abstract

This invention provides, low molecular weight, non-peptide inhibitors of matrix metalloproteinases and TNF- α converting enzyme (TACE, tumor necrosis factor- α converting enzyme) of formula (B) wherein (B) is (1), (2), or (3); P and Q are (4) or (5), provided that when P is (4), Q is (5), and vice versa; T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R¹; Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided



that no more than 2 of T, U, W, and X are nitrogen; (6) is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms which may contain 0-2 heteroatoms selected from nitrogen, oxygen, and sulfur, in addition to any heteroatoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R¹; Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteroatoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri-substituted with R¹; R¹ is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -(CH₂)_nZ, -OR², -CN, -COR², perfluoroalkyl of 1-4 carbon atoms, -CONR²R³, -S(O)₂R²-OPO(OR²)OR³, -PO(OR²)R³, -OC(O)NR²R³, -COOR², -CONR²R³, -SO₂H, -NR²R³, -NR²COR³, -NR²COOR³, -SO₂NR²R³, -NR²CONR²R³, -NR²C(-NR³)NR²R³, -SO₂NHCOR⁴, -CONHSO₂R⁴, -tetrazol-5-yl, -SO₂NHCN, -SO₂NHCONR²R³, or Z; V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteroatoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R²; R² and R³ are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V; R⁴ is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V; R⁵ is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V; n = 1-6; x = 0-2 or a pharmaceutically acceptable salt thereof.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

**THE PREPARATION AND USE OF ORTHO-SULFONAMIDO
BICYCLIC HETEROARYL HYDROXAMIC ACIDS AS MATRIX
METALLOPROTEINASE AND TACE INHIBITORS**

5 This application is a continuation-in-part of U.S. patent application SN (not obtained yet; attorney docket number 33,194-1-C1), filed on April 6, 1998, which is a continuation in part of US patent application number 08/944,188 filed on October 6,
10 1997 which claims the benefit of priority to US Provisional Application Number 60/028,505 filed October 16, 1996.

BACKGROUND

15 The present invention relates to the discovery of novel, low molecular weight, non-peptide inhibitors of matrix metalloproteinases (e.g. gelatinases, stromelysins and collagenases) and TNF- α converting enzyme (TACE, tumor necrosis factor- α converting enzyme) which are useful for the treatment of diseases in which these enzymes are implicated such as arthritis, tumor metastasis, tissue ulceration, abnormal
20 wound healing, periodontal disease, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system and HIV infection.

Matrix metalloproteinases (MMPs) are a group of enzymes that have been implicated in the pathological destruction of connective tissue and basement membranes
25 [Woessner, J.F., Jr. *FASEB J.* **1991**, *5*, 2145; Birkedal-Hansen, H.; Moore, W.G.I.; Bodden, M.K.; Windsor, L.J.; Birkedal-Hansen, B.; DeCarlo, A.; Engler, J.A. *Crit. Rev. Oral Biol. Med.* **1993**, *4*, 197; Cawston, T.E. *Pharmacol. Ther.* **1996**, *70*, 163; Powell, W.C.; Matrisian, L.M. *Cur. Top. Microbiol. and Immunol.* **1996**, *213*, 1]. These zinc containing endopeptidases consist of several subsets of enzymes including
30 collagenases, stromelysins and gelatinases. Of these classes, the gelatinases have been shown to be the MMPs most intimately involved with the growth and spread of tumors, while the collagenases have been associated with the pathogenesis of osteoarthritis [Howell, D.S.; Pelletier, J.-P. In *Arthritis and Allied Conditions*; McCarthy, D.J.; Koopman, W.J., Eds.; Lea and Febiger: Philadelphia, 1993; *12th Edition* Vol. 2, pp.
35 1723; Dean, D.D. *Sem. Arthritis Rheum.* **1991**, *20*, 2; Crawford, H.C; Matrisian, L.M. *Invasion Metast.* **1994-95**, *14*, 234; Ray, J.M.; Stetler-Stevenson, W.G. *Exp. Opin. Invest. Drugs*, **1996**, *5*, 323].

- 2 -

It is known that the level of expression of gelatinase is elevated in malignancies, and that gelatinase can degrade the basement membrane which may lead to tumor metastasis [Powell, W.C.; Matrisian, L.M. *Cur. Top. Microbiol. and Immunol.* **1996**, *213*, 1; Crawford, H.C; Matrisian, L.M. *Invasion Metast.* **1994-95**, *14*, 234; Ray, J.M.; Stetler-Stevenson, W.G. *Exp. Opin. Invest. Drugs*, **1996**, *5*, 323; Himmelstein, B.P.; Canete-Soler, R.; Bernhard, E.J.; Dilks, D.W.; Muschel, R.J. *Invasion Metast.* **1994-95**, *14*, 246; Nuovo, G.J.; MacConnell, P.B.; Simsir, A.; Valea, F.; French, D.L. *Cancer Res.* **1995**, *55*, 267-275; Walther, M.M.; Levy, A.; Hurley, K.; Venzon, D.; Linehen, W.M.; Stetler-Stevenson, W. *J. Urol.* **1995**, *153* (Suppl. 4), 403A;

10 Tokuraku, M; Sato, H.; Murakami, S.; Okada, Y.; Watanabe, Y.; Seiki, M. *Int. J. Cancer*, **1995**, *64*, 355; Himmelstein, B.; Hua, J.; Bernhard, E.; Muschel, R.J. *Proc. Am. Assoc. Cancer Res. Ann. Meet.* **1996**, *37*, 632; Ueda, Y.; Imai, K.; Tsuchiya, H.; Fujimoto, N.; Nakanishi, I.; Katsuda, S.; Seiki, M.; Okada, Y. *Am. J. Pathol.* **1996**, *148*, 611; Gress, T.M.; Mueller-Pillasch, F.; Lerch, M.M.; Friess, H.; Buechler, M.;

15 Adler, G. *Int. J. Cancer*, **1995**, *62*, 407; Kawashima, A.; Nakanishi, I.; Tsuchiya, H.; Roessner, A.; Obata, K.; Okada, Y. *Virchows Arch.*, **1994**, *424*, 547-552.]. Angiogenesis, required for the growth of solid tumors, has also recently been shown to have a gelatinase component to its pathology [Crawford, H.C; Matrisian, L.M. *Invasion Metast.* **1994-95**, *14*, 234; Ray, J.M.; Stetler-Stevenson, W.G. *Exp. Opin. Invest.*

20 *Drugs*, **1996**, *5*, 323.]. Furthermore, there is evidence to suggest that gelatinase is involved in plaque rupture associated with atherosclerosis [Dollery, C.M.; McEwan, J.R.; Henney, A.M. *Circ. Res.* **1995**, *77*, 863; Zempo, N.; Koyama, N.; Kenagy, R.D.; Lea, H.J.; Clowes, A.W. *Arterioscler. Thromb. Vasc. Biol.* **1996**, *16*, 28; Lee, R.T.; Schoen, F.J.; Loree, H.M.; Lark, M.W., Libby, P. *Arterioscler. Thromb. Vasc.*

25 *Biol.* **1996**, *16*, 1070.]. Other conditions mediated by MMPs are restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the

30 nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, periodontal disease, age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft

35 rejection.

The hypothesis that MMPs are important mediators of the tissue destruction that occurs in arthritis has long been considered, since it was first recognized that these enzymes are capable of degrading collagens and proteoglycans which are the major structural components of cartilage [Sapolsky, A.I.; Keiser, H.; Howell, D.S.; Woessner, J.F., Jr.; *J. Clin. Invest.* **1976**, *58*, 1030; Pelletier, J.-P.; Martel-Pelletier, J.; Howell, D.S.; Ghandur-Mnaymneh, L.; Enis, J.E.; Woessner, J.F., Jr., *Arthritis Rheum.* **1983**, *26*, 63.], and continues to develop as new MMPs are identified. For example, collagenase-3 (MMP-13) was cloned from breast cancer cells in 1994, and the first report that it could be involved in arthritis appeared in 1995 [Freiji, J.M.; Diez-Itza, I.; Balbin, M.; Sanchez, L.M.; Blasco, R.; Tolivia, J.; Lopez-Otin, C. *J. Biol. Chem.* **1994**, *269*, 16766; Flannery, C.R.; Sandy, J.D. **102-17**, *41st Ann. Meet. Orth. Res. Soc.* Orlando, FL. February 13-16, 1995.]. Evidence is accumulating that implicates MMP-13 in the pathogenesis of arthritis. A major structural component of articular cartilage, type II collagen, is the preferred substrate for MMP-13 and this enzyme is significantly more efficient at cleaving type II collagen than the other collagenases [Knauper, V.; Lopez-Otin, C.; Smith, B.; Knight, G.; Murphy, G. *J. Biol. Chem.*, **1996**, *271*, 1544-1550; Mitchell, P.G.; Magna, H.A.; Reeves, L.M.; Lopresti-Morrow, L.L.; Yocum, S.A.; Rosner, P.J.; Geoghegan, K.F.; Hambor, J.E. *J. Clin. Invest.* **1996**, *97*, 761.]. MMP-13 is produced by chondrocytes, and elevated levels of MMP-13 has been found in human osteoarthritic tissues [Reboul, P.; Pelletier, J.-P.; Hambor, J.; Magna, H.; Tardif, G.; Cloutier, J.-M.; Martel-Pelletier, J. *Arthritis Rheum.* **1995**, *38* (Suppl. 9), S268; Shlopov, B.V.; Mainardi, C.L.; Hasty, K.A. *Arthritis Rheum.* **1995**, *38* (Suppl. 9), S313; Reboul, P.; Pelletier, J.-P.; Tardif, G.; Cloutier, J.-M.; Martel-Pelletier, J. *J. Clin. Invest.* **1996**, *97*, 2011]. Potent inhibitors of MMPs were described over 10 years ago, but the poor bioavailability of these early peptidic, substrate mimetic MMP inhibitors precluded their evaluation in animal models of arthritis. More bioavailable, non-peptidic MMP inhibitors may be preferred for the treatment of diseases mediated by MMPs.


TNF- α converting enzyme catalyzes the formation of TNF- α from membrane bound TNF- α precursor protein. TNF- α is a pro-inflammatory cytokine that is now thought to have a role in rheumatoid arthritis, septic shock, graft rejection, insulin resistance and HIV infection in addition to its well documented antitumor properties. For example, research with anti-TNF- α antibodies and transgenic animals has demonstrated that blocking the formation of TNF- α inhibits the progression of arthritis [Rankin, E.C.; Choy, E.H.; Kassimos, D.; Kingsley, G.H.; Sopwith, A.M.; Isenberg, D.A.; Panayi,

- 4 -

G.S. *Br. J. Rheumatol.* **1995**, *34*, 334; *Pharmaprojects*, **1996**, Therapeutic Updates 17 (Oct.), 197. This observation has recently been extended to humans as well. Other conditions mediated by TNF- α are congestive heart failure, cachexia, anorexia, inflammation, fever, inflammatory disease of the central nervous system, and inflammatory bowel disease.

It is expected that small molecule inhibitors of gelatinase and TACE therefore have the potential for treating a variety of disease states. While a variety of MMP and TACE inhibitors have been identified and disclosed in the literature, the vast majority of these molecules are peptidic or peptide-like compounds that may have bioavailability and pharmacokinetic problems that would limit their clinical effectiveness. Low molecular weight, potent, long-acting, orally bioavailable inhibitors of gelatinases, collagenases and/or TACE are therefore highly desirable for the potential chronic treatment of the above mentioned disease states. Several non-peptidic, sulfur-containing hydroxamic acids have recently been disclosed and are listed below.

U. S. patents 5,455,258, 5,506,242 and 5,552,419, as well as European patent application EP606,046A1 and WIPO international publications WO96/00214 and WO97/22587 disclose non-peptide matrix metalloproteinase inhibitors of which the compound CGS27023A is representative. The discovery of this type of MMP inhibitor is further detailed by MacPherson, *et. al.* in *J. Med. Chem.*, (1997), **40**, 2525. Additional publications disclosing sulfonamide based MMP inhibitors which are variants of the sulfonamide-hydroxamate shown below, or the analogous sulfonamide-carboxylates, are European patent application EP-757984-A1 and WIPO international publications WO95/35275, WO95/35276, WO96/27583, WO97/19068 and WO97/27174.




CGS 27023A

Publications disclosing β -sulfonamide-hydroxamate MMP inhibitor analogs of CGS 27023A in which the carbon alpha to the hydroxamic acid has been joined in a


- 5 -

ring to the sulfonamide nitrogen, as shown below, include WIPO international publications WO96/33172 and WO97/20824.



5


The German patent application DE19,542,189-A1 discloses additional examples of cyclic sulfonamides as MMP inhibitors. In this case the sulfonamide-containing ring is fused to a phenyl ring to form an isoquinoline.



10

Analogs of the sulfonamide-hydroxamate MMP inhibitors in which the sulfonamide nitrogen has been replaced by a carbon atom, as shown in the general structure below, are European patent application EP-780386-A1 and WIPO international publication WO97/24117.

15



- 6 -


DESCRIPTION OF THE INVENTION

This invention provides TACE and MMP inhibitors having the formula



5

B

wherein **B** is




10



15

T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R¹;
 Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided that no more than 2 of T, U, W, and X are nitrogen;

- 7 -



is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms

which may contain 0-2 heteratoms selected from nitrogen, oxygen, and sulfur, in addition to any heteroatoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R^1 ;

Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteroatoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri- substituted with R^1 ;

R^1 is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, $-(CH_2)_nZ$, $-OR^2$, $-CN$, $-COR^2$, perfluoroalkyl of 1-4 carbon atoms, $-CONR^2R^3$, $-S(O)_xR^2$, $-OPO(OR^2)OR^3$, $-PO(OR^2)R^3$, $-OC(O)NR^2R^3$, $-COOR^2$, $-CONR^2R^3$, $-SO_3H$, $-NR^2R^3$, $-NR^2COR^3$, $-NR^2COOR^3$, $-SO_2NR^2R^3$, $-NO_2$, $-N(R^2)SO_2R^3$, $-NR^2CONR^2R^3$, $-NR^2C(=NR^3)NR^2R^3$, $-SO_2NHCOR^4$, $-CONHSO_2R^4$, -tetrazol-5-yl, $-SO_2NHCN$, $-SO_2NHCONR^2R^3$, or Z;

V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteroatoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R^2 ;

R^2 and R^3 are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

R^4 is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

R^5 is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V;

$n = 1-6$;

$x = 0-2$

- 8 -

or a pharmaceutically acceptable salt thereof.

The compounds of this invention are shown to inhibit the enzymes MMP-1, MMP-9, MMP-13 and TNF- α converting enzyme (TACE) and are therefore useful in the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection.

Pharmaceutically acceptable salts can be formed from organic and inorganic acids, for example, acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids when a compound of this invention contains a basic moiety. Salts may also be formed from organic and inorganic bases, preferably alkali metal salts, for example, sodium, lithium, or potassium, when a compound of this invention contains an acidic moiety.


Alkyl, alkenyl, alkynyl, and perfluoroalkyl include both straight chain as well as branched moieties. The definitions of alkyl, alkenyl, alkynyl, and cycloalkyl include alkyl, alkenyl, alkynyl, and cycloalkyl moieties which are unsubstituted (carbons bonded to hydrogen, or other carbons in the chain or ring) or may be mono- or poly-substituted with R¹. Halogen means bromine, chlorine, fluorine, and iodine. Preferred heteroaryl rings include pyrrole, furan, thiophene, pyridine, pyrimidine, pyridazine, pyrazine, triazole, pyrazole, imidazole, isothiazole, thiazole, isoxazole and oxazole. Preferred "heteroaryl fused to phenyl" rings include indole, isoindole, benzofuran, benzothiophene, quinoline, isoquinoline, quinoxaline, quinazoline, benzotriazole, indazole, benzimidazole, benzothiazole, benzisoxazole, and benzoxazole. The term "saturated or partially unsaturated heterocycloalkyl ring" means a saturated or partially unsaturated (but not aromatic, or fully saturated) heterocycle having 5-7 ring atoms, and containing 1-3 heteroatoms selected from N, O, or S. Preferred saturated or partially unsaturated heterocycloalkyl rings include piperidine, piperazine, morpholine, tetrahydropyran, thiomorpholine, or pyrrolidine. When a moiety contains more than one substituent with the same designation (i.e., phenyl tri-substituted with R¹) each of those substituents (R¹ in this case) may be the same or different.

The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers

- 9 -


and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry in **B**, the present invention includes such optical isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and
 5 pharmaceutically acceptable salts thereof.

Preferred compounds of this invention are those in which:



10


More preferred compounds of this invention are those in which:



- W and X are carbon; and
 15 T is nitrogen;
 U is carbon, optionally substituted with R¹
 or a pharmaceutically acceptable salt thereof.

Still more preferred compounds of this invention are those in which:
 20

- 10 -





W and X are carbon; and

T is nitrogen;

U is carbon, optionally substituted with R¹

5



is phenyl or pyrazole, each optionally mono-, di-, or tri-

substituted with R¹;

10 or a pharmaceutically acceptable salt thereof.

When R⁵ is Z, it is preferred that Z is phenyl or pyridyl, each optionally mono-, di-, or tri- substituted with R¹.






















15 It is preferred that the Z moiety bonded to the sulfur of the sulfonamide of B, is phenyl optionally mono-substituted with R¹ and R¹ is OR² or Z. When R¹ is Z it is preferred that Z is phenyl, or pyridyl, each optionally mono-, di-, or tri- substituted with R¹. When R¹ is OR², it is preferred that R² is alkyl of 1-8 carbon atoms or Z, with Z being phenyl or pyridyl, each optionally mono-, di-, or tri- substituted with R¹.

20 The compounds of this invention can be prepared according to the following schemes from commercially available starting materials or starting materials which can

- 11 -

be prepared using to literature procedures. Typical known starting materials are shown below (I-XXI). These schemes, which follow thereafter, show the preparation of representative compounds of this invention.

- 12 -



Compound I:

a) Springer, RH; Scholten, MB; O'Brien, DE, Novinson, T; Miller, JP; Robins, RK J. **Med. Chem.** (1982), 25(3), 235-42.

b) Elworthy, T.R.; Ford, A.P.D.; et.al. **J. Med. Chem.** (1997), 40(17), 2674-2687.

5 Compound II:

Masui, T; Takura, T; **JP 46043792**; JP 690307; CAN 76:59604

Compound III:

Camparini, A; Ponticelli, F; Tedeschi, P **J. Chem. Soc., Perkin Trans.1** (1982), 10, 2391-4.

10 Compound IV:

Abdalla, GM; Sowell, JW **J. Heterocycl. Chem.** (1990), 27 (5), 1201-7.

Compound V:

a) Denzel, T; Hoehn, H **J. Heterocyclic Chem.** (1977), 14, 813-817.

15 b) Al-Shaar, AHM; Chambers, RK; Gilmour, DW; Lythgoe, DJ; McClenaghan, I; Ramsden, CA **J. Chem. Soc.; Perkin Trans. I** (1992) 21, 2789-2812.

c) Elworthy, T.R.; Ford, A.P.D.; et.al. **J. Med. Chem.** (1997), 40(17), 2674-2687.

Compound VI:

a) Forbes, IT; Johnson, CN; Jones, GE; Loudon, J; Nicholass, JM **J. Med. Chem** (1990) 2640- 2645.

20 b) Kan, MA; Guarconi, AE **J. Heterocyclic Chem** (1977) 14, 807-812.

Compound VII:

a) Forbes, IT; Johnson, CN; Jones, GE; Loudon, J; Nicholass, JM **J. Med. Chem** (1990) 2640- 2645.

b) Kan, MA; Guarconi, AE **J. Heterocyclic Chem** (1977) 14, 807-812.

25 Compound VIII:

Richardson, TO; Neale, N; Carwell, N **J. Heterocyclic. Chem.** (1995), 32, 359-361.

Baker, JM; Huddleston, PR; Keenan, GJ **J. Chem Research Miniprint**, (1982) 6, 1726-1746.

30 Compound IX:

a) Forbes, IT; Johnson, CN; Jones, GE; Loudon, J; Nicholass, JM **J. Med. Chem** (1990) 2640- 2645.

b) Kan, MA; Guarconi, AE **J. Heterocyclic Chem** (1977) 14, 807-812.

Compounds X, XI and XII:

35 Elworthy, T.R.; Ford, A.P.D.; et.al. **J. Med. Chem.** (1997), 40(17), 2674-2687.

- 14 -

Compound XIII:

Heterocycles, (1997), 45, 980.

Compound XIV:

5 Yokoyama, Naokata. Eur. Pat. Appl., 61 pp. CODEN: EPXXDW. **EP 115469 A1**
840808.

Compound XV:

Mendes, Etienne; Vernieres, Jean Claude; Simiand, Jacques Edouard; Keane, Peter Eugene. Eur. Pat. Appl., 12 pp. CODEN: EPXXDW. **EP 346207 A1** 891213.

Compound XVI:

10 Mendes, Etienne; Vernieres, Jean Claude; Simiand, Jacques Edouard; Keane, Peter Eugene. Eur. Pat. Appl., 12 pp. CODEN: EPXXDW. **EP 346207 A1** 891213.

Compound XVII:

Morita, Yoshiharu; Wagatsuma, Kazuo. Japan. Kokai, 4 pp. CODEN: JKXXAF. **JP 50058094** 750520 Showa.

15 Compounds XVIII and XIX:

Armitage, Bernard John; Leslie, Bruce William; Miller, Thomas Kerr; Morley, Christopher. PCT Int. Appl., 110 pp. CODEN: PIXXD2. **WO 9500511 A1** 950105.

Compound XX:


20 Minami, S.; Matsumoto, J.; Kawaguchi, K.; Mishio, S.; Shimizu, M.; Takase, Y.; Nakamura, S. (Dainippon Pharmaceutical Co., Ltd., Japan) Japan. Kokai, 3pp. CODEN: JKXXAF. **JP 50014697** 750215 Showa.

Compound XXI:

25 Kihara, N.; Tan, H.; Takei, M.; Ishihara, T. (Mitsui Pechochemical Industries, Ltd., Japan; Suntory, Ltd.) Jpn. Kokai Tokyo Koho, 11pp. CODEN: JKXXAF. **JP 62221686 A2** 870929 Showa.

The compounds of this invention can be prepared using conventional techniques known to those skilled in the art of organic synthesis. The following scheme (Scheme I)
30 illustrates the reaction sequence employed. In the schemes which follow, the moiety A is defined as the bicyclic heteroaryl moiety of B, as shown immediately below:

- 15 -




For purposes of illustration only, wherein the bicyclic heteroaryl group A shown is a quinoline, 4-chloro-7-trifluoromethylquinoline-3-carboxylic acid ethyl ester, prepared from the corresponding aniline, is reacted with *N*-benzyl-*p*-methoxybenzenesulfonamide, wherein Z is *p*-methoxybenzene, to provide the requisite *N,N*-disubstituted sulfonamido-ester which is then converted into the corresponding hydroxamic acid in two steps.

Alternatively, the 4-chloroquinoline carboxylic acid ester could be first reacted with R^7-NH_2 and the resulting 4-(R^7 -amino)quinoline carboxylic acid ester then reacted with the appropriate $Z-SO_2-Cl$. Hydrolysis of the ester and reaction with hydroxylamine hydro-chloride would then give the invention compound.

- 16 -

Scheme I.




5

Functionalization of the quinoline ring via a palladium catalyzed Heck coupling between the iodoquinoline and tributylvinyltin is shown in Scheme II. α,β -Unsaturated esters and amides can be coupled to the haloquinoline via Heck reactions. A variety of other trialkyltin reagents are readily available and may be similarly used. Boronic acids, commercially available or readily prepared, may also be coupled to the iodoquinoline

10 using the Suzuki reaction.

- 17 -


Scheme II.



5 Functionalization of haloquinolines may also be accomplished via palladium catalyzed couplings of alkynes, as illustrated in Scheme III. Hydrogenation of the alkynes accesses the olefins and alkanes as well.

- 18 -


Scheme III.



Schemes IV and V illustrate two methods for incorporating amino groups into the substituent attached to the sulfonamide nitrogen of the compounds of the invention. Thus, in Scheme IV the NH-sulfonamide is alkylated with propargyl bromide to provide the propargyl sulfonamide. This alkyne is reacted with paraformaldehyde in the presence of a primary or secondary amine and cuprous chloride to give the propargyl amine which is converted, as before, to the desired hydroxamic acid.

10


Scheme IV.



- 5 In Scheme V, selective hydrolysis of the ester of the p-carboethoxybenzyl sulfonamide group provides a mono-carboxylic acid. This acid may be converted into an amide (not shown), followed by conversion of the second ester, A-CO₂R, into the corresponding hydroxamate, or reduced to the corresponding alcohol with diborane. The alcohol may be converted into the analogous amine via the benzylic bromide, followed by conversion of the the ester, A-CO₂R, into the corresponding hydroxamate.
- 10


- 20 -

Scheme V.



- 5 Methods for synthesizing variations of substituents on the sulfonyl aryl group are shown in Schemes VI through VIII. As shown in Scheme VI, biaryl sulfonyl groups are synthesized by Suzuki couplings on a bromo-substituted benzene sulfonamide. The starting bromo-substituted benzene sulfonamide is synthesized from the commercially available bromobenzenesulfonyl chloride and the amino-acid or amino-ester, $\text{H}_2\text{N-A-CO}_2\text{R}$, followed by alkylation of the resulting NH-sulfonamide. Alternatively, the
- 10 bromo aryl sulfonamide is converted into the corresponding boronic acid by the method of Ishiyama, et.al. [J. Org. Chem. (1995), 60, 7508] followed by coupling with an appropriate aryl halide.


Scheme VI.



- 5 Methods for synthesizing sulfonyl aryl ethers are shown in Schemes VII through IX. In Scheme VII biaryl ethers, or aryl heteroaryl ethers, are synthesized starting from the known sulfonyl chlorides (see for example: Zook SE; Dagnino, R; Deason, ME, Bender, SL; Melnick, MJ WO 97/20824).


- 22 -

Scheme VII.



Alternatively, the biaryl ethers may be prepared from the corresponding boronic
 5 acids or via the sulfonyl phenols as shown in Scheme VIII.


Scheme VIII.



- 5 Aryl ethers may also be prepared via displacement of the fluorine from a para-fluorobenzene sulfonamide, as shown in Scheme IX. Aryl or alkyl ethers may be prepared in this manner.


- 24 -

Scheme IX.




- 5 Scheme X illustrates the synthesis of pyrazolopyridines, isoxazolopyridines, and isothiazolopyridines of the invention. Thus, an aminopyrazole, aminoisoxazole or aminoisothiazole is condensed with ethoxymethylene malonate to provide the intermediate, B. This compound is converted into the pyrazolopyridine, isoxazolopyridine, or isothiazolopyridine, C, by heating at 240°C. Compound C is then
- 10 converted into the chloro-ester, D, via reaction with phosphorus oxychloride. Displacement of the chloro substituent with a sulfonamide then gives compound E. Hydrolysis of the ester and conversion of the carboxylate into the hydroxamate then gives compound G. Salts of the invention compounds can be prepared according to standard procedures.

Scheme X



- 26 -

- 5 Pyrazolo[1,5-b]pyrimidines of the invention are prepared according to scheme XI using reactions as described for scheme X.



10

The following specific examples illustrate the preparation of representative compounds of this invention. The starting materials, intermediates, and reagents are either commercially available or can be readily prepared following standard literature procedures by one skilled in the art of organic synthesis.

15

- 27 -

Example 1

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester

5 To a solution of 1.85g (6.67 mmol) of N-benzyl 4-methoxyphenylsulphonamide in 15mL of DMF was added, in one portion, 0.267g (6.67 mmol) of 60% sodium hydride and the resulting mixture was stirred at room temperature under nitrogen for 15 min. Ethyl 4-chloro-7-trifluoromethyl-3-quinolinecarboxylate (2.02g, 6.67 mmol) was then added to the solution in one portion and the resulting mixture was heated at 85 °C
10 for 24h. The reaction mixture was then cooled to room temperature, poured into a mixture of water (300 mL) and HCl (1N, aqueous, 100 mL) and extracted with ethyl acetate (2x100 mL). The combined organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo. The residue was then chromatographed on silica gel eluting with 15%-50% ethyl acetate/ hexane to give 3.11g (88%) of the desired product.
15 Electrospray Mass Spec 545.1 (M+H).

Example 2

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-8-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester

20 In the same manner as described in Example 1, 1.012g (3.34 mmol) of ethyl 4-chloro-8-trifluoromethyl-3-quinolinecarboxylate provided 1.509g (83%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 545.1 (M+H).

Example 3

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]- 6-bromo-quinoline-3-carboxylic acid ethyl ester

25 In the same manner as described in Example 1, 0.848g (2.70 mmol) of ethyl 6-bromo-4-chloro-3-quinolinecarboxylate provided 1.418g (95%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 557.1 (M+H).
30

- 28 -

Example 4

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-bromo-quinoline-3-carboxylic acid ethyl ester

5 In the same manner as described in Example 1, 0.777g (2.47 mmol) of ethyl 7-bromo-4-chloro-3-quinolinecarboxylate provided 1.169g (85%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 557.1 (M+H).

Example 5

10 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester**

In the same manner as described in Example 1, 1.216g (4.02 mmol) of ethyl 4-chloro-6-trifluoromethyl-3-quinolinecarboxylate provided 2.171g (99%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 545.0 (M+H).

15 Example 6

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid

20 To a solution of 1.065g (2.00 mmol) of the product from Example 1 in 4mL of methanol/THF (1:1) was added 2mL of 1N sodium hydroxide solution and the resulting mixture was stirred at 25 °C for 18h. The reaction was then acidified with 1N HCl and extracted with ethyl acetate (200 mL). The organic layer was washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The resulting residue was triturated with ethyl acetate/hexane (1:9) and filtered to provide 828 mg (82%) of the desired carboxylic acid as a white solid. Electrospray Mass Spec 517.1 (M+H)

25

Example 7

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-8-trifluoromethyl-quinoline-3-carboxylic acid

30 In the same manner as described in Example 6, 1.255g (2.64 mmol) of the product from Example 2 provided 0.988g (83%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 517.1 (M+H).

- 29 -

Example 8

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-bromo-quinoline-3-carboxylic acid

5 In the same manner as described in Example 6, 1.198g (2.16 mmol) of the product from Example 3 provided 0.921g (81%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 529.0 (M+H).

Example 9

10 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]- 7-bromo-quinoline-3-carboxylic acid**

In the same manner as described in Example 6, 0.969g (1.74 mmol) of the product from Example 4 provided 0.804g (87%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 529.0 (M+H).

15

Example 10

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-trifluoromethyl-quinoline-3-carboxylic acid

20 In the same manner as described in Example 6, 2.043g (3.75 mmol) of the product from Example 5 provided 1.82g (88%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 515.0 (M-H).

Example 11

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

25 To a solution of 0.636g (1.26 mmol) of the product from Example 6 in 12.5 mL of dichloromethane was added 0.05 mL of DMF followed by 1.26 mL (2.52 mmol) of 2 M oxalyl chloride and the resulting reaction mixture was stirred at room temperature for 1h.

30 In a separate flask, 2.6 mL (19 mmol) of triethylamine was added to a 0°C mixture of 350 mg (13 mmol) of hydroxylamine hydrochloride in 14 mL of THF and 3.5 mL of water. After this mixture had been stirred for 15min at 0 °C, the acid chloride solution was added to it in one portion and the resulting solution was allowed to warm to room temperature and stirred for another 4h. Water was then added to the reaction flask and 0.488g (75%) product was collected via filtration. Electrospray Mass Spec 532.1
35 (M+H)

- 30 -

Example 12

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-8-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

5 In the same manner as described in Example 11, 0.444g (3.75 mmol) of the product from Example 7 provided 0.143g (31%) of the desired quinoline hydroxamic acid as a cream colored solid. Electrospray Mass Spec 532.1 (M+H).

Example 13

10 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]- 6-bromo-quinoline-3-carboxylic acid hydroxyamide**

In the same manner as described in Example 11, 0.527g (1.00 mmol) of the product from Example 8 provided 0.367g (68%) of the desired quinoline hydroxamic acid as a off-white solid. Electrospray Mass Spec 541.9 (M+H).

15

Example 14

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]- 7-bromo-quinoline-3-carboxylic acid hydroxyamide

20 In the same manner as described in Example 11, 0.527g (1.00 mmol) of the product from Example 9 provided 0.280g (52%) of the desired quinoline hydroxamic acid as a white solid. Electrospray Mass Spec 541.9 (M+H).

Example 15

25 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide**

In the same manner as described in Example 11, 0.527 g (1.06 mmol) of the product from Example 10 provided 0.435g (77%) of the desired quinoline hydroxamic acid as a cream colored solid. Electrospray Mass Spec 532.1 (M+H).

30

Example 16

4-[(4-Methoxybenzenesulfonyl)-pyridin-3-ylmethylamino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

35 Following the procedure of example 1 and substituting N-(3-pyridinylmethyl)-4-methoxybenzenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide, the intermediate 4-[(4-methoxybenzenesulfonyl)-pyridin-3-ylmethylamino]-7-

trifluoromethyl-quinoline-3-carboxylic acid ethyl ester is obtained. Following the procedures of example 6 and 11, the title product is obtained. Electrospray Mass Spec 533.0 (M+H).

5

Example 17

4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-t-butyl-quinoline-3-carboxylic acid hydroxyamide

In the same manner as described in Example 1, 1.167g (4.00 mmol) of ethyl 4-chloro-8-butyl-3-quinolinecarboxylate provided 1.413g (66%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 533.3 (M+H).

10

In the same manner as described in Example 6, 1.065g (2.00 mmol) of the ester provided 0.478g (47%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 503.3 (M-H).

Following the procedures of example, the title compound is obtained from the carboxylic acid. Electrospray Mass Spec. 520.3 (M+H).

15

Example 18

4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-methyl-quinoline-3-carboxylic acid hydroxyamide

20

In the same manner as described in Example 1, 1.00g (4.00 mmol) of ethyl 4-chloro-8-methyl-3-quinolinecarboxylate provided 0.531g (27%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 491.3 (M+H).

In the same manner as described in Example 6, 0.470g (0.851 mmol) of the ester provided 0.160g (41%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 461.3 (M-H).

25

Following the procedure of example 11, the title compound is obtained from the carboxylic acid. Electrospray Mass Spec. 478.3 (M+H).

Example 19

30

4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-ethyl-quinoline-3-carboxylic acid hydroxyamide

In the same manner as described in Example 1, 1.055g (4.00 mmol) of ethyl 4-chloro-8-ethyl-3-quinolinecarboxylate provided 0.670g (33%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 505.3 (M+H).

- 32 -

In the same manner as described in Example 6, 0.615g (1.22 mmol) of the product from Example 7 provided 0.353g (60%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 475.3 (M-H).

Following the procedure of example 11, the title compound is obtained from the
5 carboxylic acid. Electrospray Mass Spec. 492.3 (M+H).

Example 20

4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-(1-methylethyl)- quinoline-3-carboxylic acid hydroxyamide

10 In the same manner as described in Example 1, 1.111g (4.00 mmol) of ethyl 4-chloro-8-isopropyl-3-quinolinecarboxylate provided 0.754g (36%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 519.3 (M+H).

In the same manner as described in Example 6, 0.686g (0.127 mmol) of the ester
15 provided 0.532g (82%) of the desired quinoline acid as a white solid. Electrospray Mass Spec 489.2 (M-H).

In the same manner as described in Example 11, 0.440g (0.897 mmol) of the hydroxamic acid provided 0.270 g (60%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 506.3 (M+H).

20 Example 21

4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]- 8-iodo-quinoline-3- carboxylic acid ethyl ester

In the same manner as described in Example 1 and substituting N-ethyl-4-methoxybenzenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide, 1.076g
25 (5.00 mmol) of ethyl 8-iodo-4-chloro-3-quinolinecarboxylate provided 2.438g (4.51 mmol, 90%) of the desired quinoline ester as a white solid. Electrospray Mass Spec 541.0 (M+H).

Example 22

4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-8-vinyl-quinoline-3- carboxylic acid ethyl ester

The product from example 21 (2.438g, 4.51mmol) in 150 mL DMF was added tributylvinyltin (1.43g, 4.51 mmol), tetrakis(triphenylphosphine)palladium(0) (520mg, 10%), cuprous iodide (171mg, 20%), and 5 mL triethylamine. The mixture was stirred
35 under N₂ and heated at 85°C for 18 hours. The it was poured into a mixture (1:1) of 400

- 33 -

mL saturated sodium bicarbonate and saturated ammonium chloride and extracted with ethyl acetate (2x200 mL). The combined organic layers were dried over magnesium sulfate, filtered and concentrated on a rotary evaporator. The residue was column chromatographed using 300 mL silica gel and gradient elution with hexane/ethyl acetate (100-0%). This provided 1.706g (3.88 mmol, 86%) of the desired quinoline ester. Electropray Mass Spec 441.1 (M+H).

Example 23

4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethynyl-quinoline-3-carboxylic acid ethyl ester

Combining the procedures of examples 1 and 22, and substituting phenylacetylene for vinyltin, N-ethyl-4-methoxybenzenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide, the intermediate 4-[ethyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethynyl-quinoline-3-carboxylic acid ethyl ester is obtained from ethyl 4-chloro-3-quinolinecarboxylate. Electropray Mass Spec 515.3 (M+H).

Example 24

4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-8-vinyl-quinoline-3-carboxylic acid

In the same manner as described in Example 6, 1.593g (3.62 mmol) of the product from Example 22 provided 1.333g (89%) of the desired quinoline acid as a white solid. Electropray Mass Spec 411.1 (M-H).

Example 25

4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethynyl-quinoline-3-carboxylic acid

In the same manner as described in Example 6, the title compound was synthesized from the product of example 23. Electropray Mass Spec 485.3 (M-H).

Example 26

4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-nitro-quinoline-3-carboxylic acid

In the same manner as described in Example 1 and 6, 5.613g (20.0 mmol) ethyl 4-chloro-6-nitro-3-quinolinecarboxylate provided 2.676g (27% for two steps) of the title compound as a white solid. Electropray Mass Spec 492.3 (M-H).

- 34 -

Example 27

4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]- 8-bromo-quinoline-3-carboxylic acid

5 Combining the procedures of example 1 and 6, and substituting N-methyl-4-methoxybenzenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide, the intermediate 8-bromo-4-[methyl-(4-methoxy-benzenesulfonyl)-amino]-quinoline-3-carboxylic acid is obtained. Electrospray Mass Spec 449.2 (M-H).

Example 28

10 **4-{Methyl-(4-(pyridin-4-yloxy)-benzenesulfonyl)-amino}-6-iodo-quinoline-3-carboxylic acid**

Combining the procedures of example 1 and 6, and substituting N-methyl-4-(pyridin-4-yloxy)-benzenesulfonamide the intermediate 6-iodo-4-{methyl-(4-(pyridin-4-yloxy)-benzenesulfonyl)-amino}-quinoline-3-carboxylic acid is obtained from ethyl 6-iodo-4-chloro-3-quinolinecarboxylate. Electrospray Mass Spec 559.9 (M-H).

15

Example 29

20 **4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-8-vinyl-quinoline-3-carboxylic acid hydroxyamide**

In the same manner as described in Example 11, 0.484g (1.17 mmol) of the product from Example 24 provided 0.360g (72%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 428.0 (M+H).

Example 30

25 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-nitro-quinoline-3-carboxylic acid hydroxyamide**

In the same manner as described in Example 11, 0.825g (1.67 mmol) of the product from Example 26 provided 0.227g (0.446 mmol, 26%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 509.0 (M+H).

30

- 35 -

Example 31

4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]- 8-bromo-quinoline-3-carboxylic acid hydroxyamide

In the same manner as described in Example 11, 0.664g (1.47 mmol) of the product from Example 27 provided 0.145g (0.311 mmol, 21%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 468.0 (M+H).

Example 32

4-{Methyl-[4-(pyridin-4-yloxy)-benzenesulfonyl]-amino}-6-iodo-quinoline-3-carboxylic acid hydroxyamide

To a 0 °C solution of of 4.5 mL oxalyl chloride (0.90 mmol, 2M in dichloromethane) was added dropwise 0.69 mL of DMF. The resulting solid was kept at 0 °C for another 15 minutes and followed by addition of 2.50g (4.46 mmol) of the product from Example 28 in 50 mL DMF. The mixture was stirred for 1 hour at room temperature and then kept at 0 °C for an additional 15 minutes. An aqueous solution of hydroxylamine (5mL, 50%) was then added all at once to the above solution and the mixture was stirred at room temperature for 3 hours. The mixture was next poured into 300mL water and extracted with dichloromethane (4x100mL). The combined organic layers were washed with brine (300mL) and dried over magnesium sulfate. After filtration and concentration on a rotary evaporator the residue was column chromatographed using gradient methanol in ethyl acetate (20-100%) and it provided 1.36g (2.36mmol, 53%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 576.9 (M+H).

Example 33

4-{Methyl-(4-(pyridin-4-yloxy)-benzenesulfonyl)-amino}-6-iodo-quinoline-3-carboxylic acid hydroxyamide hydrochloride

The product from example 32 (0.952 g, 1.65 mmol) was dissolved in 100 mL methanol in a Parr reactor. Degussa catalyst (10% Pd-C, 200mg) was next added under N₂. The mixture was then hydrogenated (35 psi) for one hour at room temperature. The mixture was then filtered through a pad of celite and concentrated on a rotary evaporator. The residue was chromatographed with methanol and ethyl acetate (5-35%). The product obtained was next dissolved in methanol and anhydrous hydrochloride was bubbled into the solution for 5 minutes. Removal of the solvent through rotary

- 36 -

evaporation and vacuum pump gave 0.707g (1.45 mmol, 88%) product. Electrospray Mass Spec. 450.9 (M+H).

Example 34

5 **4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethynyl-quinoline-3-carboxylic acid hydroxyamide**

In the same manner as described in Example 11, 2.432g (5.00 mmol) of the product from Example 25 provided 2.159g (86%) of the desired quinoline hydroxamic acid. Electrospray Mass Spec. 502.1 (M+H).

10

Example 35

15 **4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethyl-quinoline-3-carboxylic acid hydroxyamide**

The product from example 34 (0.82g, 1.64 mmol) was dissolved in 50 mL methanol in a Parr reactor. Degussa Catalyst (10% Pd-C, 200mg) was next added under N₂. The mixture was hydrogenated (45 psi) for one hour at room temperature. The mixture was then filtered through a pad of celite and concentrated on a rotary evaporator. This gave 0.76g (1.50 mmol, 92%) product. Electrospray Mass Spec. 506.0 (M+H).

20

Example 36

25 **4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]- 8-methoxy-quinoline-3-carboxylic acid hydroxyamide**

Following the procedure of Example 16 and starting with ethyl 4-chloro-8-methoxy-3-quinolinecarboxylate the title compound was obtained as a yellow powder. Electrospray Mass Spec. 495.3 (M+H).

Example 37

30 **4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]- 8-bromo-quinoline-3-carboxylic acid hydroxyamide**

Following the procedure of Example 16 and starting with ethyl 4-chloro-8-bromo-3-quinolinecarboxylate the title compound was obtained as a white powder. Electrospray Mass Spec. 543.2 (M+H).

- 37 -

Example 38

4-[(4-methoxy-benzenesulfonyl)-pyridin-3-ylmethyl amino]-8-Benzyl-quinoline-3-carboxylic acid hydroxyamide

Following the procedure of Example 16 and starting with ethyl 4-chloro-8-benzyl-3-quinolinecarboxylate the title compound was obtained as a beige powder. Electropray Mass Spec. 555.4 (M+H).

Example 39

4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]- 8-iodo-quinoline-3-carboxylic acid hydroxyamide

Following the procedure of Example 16 and starting with ethyl 4-chloro-8-iodo-3-quinolinecarboxylate the title compound was obtained as a yellow powder. Electropray Mass Spec. 590.8 (M+H).

Example 40

4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-phenyl-quinoline-3-carboxylic acid hydroxyamide

Following the procedure of Example 16 and starting with ethyl 4-chloro-8-phenyl-3-quinolinecarboxylate the title compound was obtained as a beige powder. Electropray Mass Spec. 541.4 (M+H).

Example 41

4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-thiophen-2-yl-quinoline-3-carboxylic acid hydroxyamide

Combining the procedures of Examples 22, 6 and 11 and starting with 4-[(4-methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]- 8-bromo-quinoline-3-carboxylic acid ethyl ester and 2-tributylstannylthiophene the title compound was obtained as a yellow powder. Electropray Mass Spec. 545.0 (M+H).

Example 42

4-[(Biphenyl-4-sulfonyl)-pyridin-3-ylmethyl-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

Following the procedure of Example 1 and substituting N-(3-pyridinylmethyl)-4-bromobenzenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide, the

- 38 -

intermediate 4-[(4-bromobenzenesulfonyl)-pyridin-3-ylmethylamino]-7-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester is obtained.

To 8.5 mL of degassed ethylene glycol dimethyl ether, was added 500 mg (0.85 mmol) of the ester, 110 mg (0.93 mmol) of phenylboronic acid, 80 mg (0.07 mmol) of tetrakis(triphenylphosphine)palladium and 1.1 ml (2.2 mmol) of 2M aqueous Na₂CO₃ and the mixture was heated to reflux under nitrogen for 36 hr. The reaction was cooled to room temperature, diluted with ethyl acetate, washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo to give 4-[(biphenyl-4-sulfonyl)-pyridin-3-ylmethyl-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester.

This ester was converted to the title compound (off-white powder) as described in Examples 6 and 11. Electrospray Mass Spec. 579.1 (M+H).

Example 43

4-[(Octane-1-sulfonyl)-pyridin-3-ylmethyl-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

Combining the procedures of Examples 1, 6 and 11 and substituting N-(3-pyridinylmethyl)-octanesulfonamide for N-benzyl-4-methoxybenzenesulfonamide the title compound was obtained as a yellow solid. Electrospray Mass Spec. 539.5 (M+H).

Example 44

4-[Pyridin-3-ylmethyl-(toluene-4-sulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide

Combining the procedures of Examples 1, 6 and 11 and substituting N-(3-pyridinylmethyl)-toluenesulfonamide for N-benzyl-4-methoxybenzenesulfonamide the title compound was obtained as a white powder. Electrospray Mass Spec. 517.1 (M+H).

Example 45

Diethyl[(1-phenyl-5-pyrazolyl) amino] methylene malonate

A mixture of 15.9 g. (0.10 mole) of 1-phenyl-5-aminopyrazole and 21.6 g. (0.10 mole) of diethyl ethoxymethylenemalonate was heated at 115-120° in an oil bath for 2 hours. After cooling, the crystalline mass was recrystallized from hot hexane containing 1% of ethanol. Cooling to room temperature and filtering gave 24.8 g. (75%) of off-white crystals, m.p. 96-97°C.

Example 46

Ethyl 4-hydroxy-1-phenyl-1H-pyrazolo [3,4-b] pyridine-5-carboxylate

A mixture of 18.1 g. (0.055 mole) of diethyl CCOC(=O)C(=O)OCC [(1-phenyl-5-pyrazolyl)amino]methylene} malonate and 150 ml of diethyl phthalate was heated at 240-250° for 1 hour. The mixture was chilled and diluted with hexane. Chilling and filtering gave crystals which were washed with hexane and with hexane-ethanol (1:1) to give 11 g. (70%) of off white crystals m.p. 149-150°C. From a similar small scale run 1.75 g. was recrystallized from 110 ml. of ethanol to give 1.58 g. of off white crystals, m.p. 149-150° C.

Example 47

Ethyl 4-chloro-1-phenyl-1H-pyrazolo [3,4-b]pyridine-5-carboxylate

A mixture of 5.76 g (20.33 mmol) of ethyl 4-hydroxy-1-phenyl-1H-pyrazolo[3,4-b] pyridine-5- carboxylate and 15.58 g of phosphorus oxychloride was refluxed 1.5 hr, chilled and poured slowly onto crushed ice. The mixture was filtered and the solid washed with ice-water and dried to give 6.0 g of solid, m.p. 89-91° C.

Example 48

Ethyl 4-chloro-1,3-dimethyl-1H-pyrazolo [3,4-b]pyridine-5-carboxylate

Following the procedures of Examples 45, 46 and 47, starting from 1,3-dimethyl-5-aminopyrazole, the chloro-ester is prepared. m.p. 89-90° C.

Example 49

Ethyl 4-[benzyl-(4-methoxybenzenesulfonyl)amino]- 1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylate

To a solution of 1.16 g (4.2 mmol) of benzyl-(4-methoxybenzenesulfonyl)amine in 6 ml of anhydrous 1-methyl-2-pyrrolidinone was added 0.168 g (4.2 mmol) of sodium hydride (60% in oil) and the mixture stirred at room temperature until gas evolution ceased. The preceding mixture was added to mixture of 1.01 g (4 mmol) of ethyl 4-chloro-1,3- dimethylpyrazolo[3,4-b]pyridine -5-carboxylate in 2 ml of 1-methyl-2-pyrrolidinone.

The mixture was heated in an oil bath at 50°C overnight and then was heated in an oil bath at 100° C for 1.5 days. The mixture was poured into 800 ml of water and extracted with ethyl acetate. The extract was washed with water, 2N citric acid, water,

- 40 -

brine and dried (Na_2SO_4). The solvent was removed and the residue chromatographed on silica gel with hexane-ethyl acetate (2:1) as eluent to give 0.64 g of product as a solid, mp 170-172°. From a larger scale run of 5.07 g (0.02 mmol) of ethyl 4-chloro-1,3-dimethylpyrazolo [3,4-b]pyridine-5-carboxylate and 8.0 g (0.0289 mmol) of benzyl -(4-methoxybenzenesulfonyl) amine (as sodium anion) in 30 ml of 1-methyl-2-pyrrolidinone heated at 90° C for 3 days there was obtained 3.65 g of product.

Example 50

4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo [3,4-b] pyridine-5-carboxylic acid.

A mixture of 0.48 g (0.97 mmol) of ethyl 4-[benzyl-(4-methoxybenzenesulfonyl) amino] -1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate and 0.29 ml of 10N NaOH in 4 ml of tetrahydrofuran-methanol (1:1) was heated in an oil bath at 70° C for 2 hours and the solvent removed under vacuum. The residue was dissolved in 20 ml of H_2O and the solution extracted with 10 ml of diethyl ether. To the aqueous layer was added 2N citric acid (pH 4-5) and the precipitated solid filtered and washed with H_2O to give a white solid which was dried under vacuum overnight to give crystals, mp 165-167°C.

Example 51

4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, potassium salt.

A mixture of 3.60 g (7.28 mmol) of ethyl 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate and 0.44 g (7.84 mmol) of potassium hydroxide (pellet) in 15 ml of methanol- water (1:1) was refluxed overnight. An additional 40 mg of potassium hydroxide was added and the mixture refluxed for 4 hours (all the solid dissolved). The solvent was removed under vacuum and toluene added and removed under vacuum. The residue was triturated with ethyl acetate, filtered and the solid washed with ethyl acetate to give 3.8 g of product as a white solid.

Example 52

4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, hydroxyamide

To a chilled solution of 1 ml (2 mmol) of oxalyl chloride in 8 ml of CH_2Cl_2 was added dropwise 0.154 ml (2 mmol) of N, N-dimethylformamide and the solution stirred 15 min. To the preceding chilled solution was added 0.504 g (1 mmol) of 4-[benzyl-(4-methoxybenzenesulfonyl) amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, potassium salt and the mixture stirred under nitrogen for 2 hrs at room temperature (solution A). A solution of 0.278 g (4 mmol) of hydroxylamine hydrochloride and 0.834 ml (6 mmol) of triethylamine in 5 ml of H_2O -tetrahydrofuran (1:4) was chilled at in an ice bath for 20 min. and to this solution was added dropwise the chilled solution of A. The mixture was allowed to warm to room temperature and was stirred overnight. The solvent was removed and the residue extracted with CH_2Cl_2 . The CH_2Cl_2 extract was washed with 2N citric acid, H_2O , 1N NaHCO_3 , H_2O , brine and dried (Na_2SO_4). The solvent was removed to give 0.53g of solid. Trituration with ethyl acetate gave 0.278 g of white solid, mp 184-186°C.

Example 53

Ethyl 4-[(4-methoxybenzenesulfonyl)pyridin -3-ylmethylamino] -1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate

To a solution of 1.39 g (5 mmol) of (4-methoxybenzenesulfonyl)(3-pyridinylmethyl) amine in 4 ml of anhydrous 1-methyl-2-pyrrolidinone was added 0.2 g (5 mmol) of sodium hydride (60% in oil) and the mixture stirred at room temperature until gas evolution ceased. To this mixture was added 1.15 g (4.54 mmol) of ethyl 4-chloro-1,3-dimethylpyrazolo[3,4-b]pyridine-5-carboxylate and 2 ml of anhydrous 1-methyl-2- pyrrolidinone. The mixture was stirred in a sealed tube under nitrogen in an oil bath at 90°C for 3 days. The mixture was cooled, poured into water and extracted with ethyl acetate. The extract was washed with H_2O , brine and dried (Na_2SO_4). The solution was filtered through a thin pad of hydrous magnesum silicate and the filter pad washed with ethyl acetate. The filtrate was concentrated to dryness under vacuum to give 1.3 g of solid. Chromatography on silica gel with ethyl acetate as solvent gave 0.35 g of product as a solid, mp 152-154°C.

- 42 -

Example 54

4-[(4-Methoxybenzenesulfonyl) pyridin -3-ylmethylamino] -1,3-dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid

A mixture of 1.34 g (2.7 mmol) of ethyl 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate, 2.97 ml of 1N potassium hydroxide in 7.8 ml of ethanol and 4.83 ml of water was refluxed for 20 hr. Another 0.54 ml of 1N potassium hydroxide was added and the mixture refluxed 4 hrs. The solvent was removed under vacuum and toluene added and removed under vacuum. The residue was dissolved in water (20ml) and extracted with ethyl acetate. The aqueous layer was acidified with 2 N citric acid and the precipitated solid filtered off and washed with water. The solid was dried under vacuum to give 0.98 g of solid, mp 256-258°C.

Example 55

4-[(4-Methoxybenzenesulfonyl) pyridin -3-ylmethylamino] -1,3-dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, potassium salt

A mixture of 0.34 g (0.68 mmol) of ethyl 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate and 0.748 ml of 1 N potassium hydroxide in 4 ml of ethanol- water (1:1) was refluxed for 24 hr. The solvent was removed under vacuum and to the residue was added toluene. The solvent was removed under vacuum to remove the water and the residue triturated with ethyl acetate to give the product as a solid, mp 160-167° C.

Example 60

4-[(4-Methoxybenzenesulfonyl)pyridin -3-ylmethylamino] -1,3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

A 1.5 g (2.459 mmol) sample of 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid was dissolved in 2.70 ml of 1N KOH. The water was removed by repeated additions and removal of toluene under vacuum to give 1.34 g of solid (potassium salt of the acid). A solution of 2.65 ml (5.3 mmol) of oxalyl chloride was cooled in an ice bath and 0.389 ml of N,N-dimethylformamide added dropwise. After 5 min. the 1.34 g of the previously prepared potassium salt was added and the mixture stirred for 10 min. in an

- 43 -

ice bath and then allowed to warm to room temperature (mixture A). A mixture of 0.737 g (10.6 mmol) of hydroxylamine hydrochloride and 2.21 ml (15.9 mmol) of triethylamine in 9.39 ml of tetrahydrofuran and 2.45 ml of water was chilled in an ice bath (mixture B). The mixture A was chilled in an ice bath and added to the chilled and stirred mixture B. The mixture of A and B was stirred at 0°C for 10 min and allowed to warm to room temperature and stir overnight. The solvent was removed under vacuum and the residue diluted with H₂O, acidified with 2 N citric acid and extracted with two 30-ml portions of CH₂Cl₂. The aqueous layer was neutralized with solid NaHCO₃ to bring the pH to 7. The solid which precipitated was filtered and washed with H₂O to give 0.610 g of product as a solid, mp. 202-204°C. The CH₂Cl₂ extract was extracted with 2 N citric acid and the aqueous layer neutralized with solid NaHCO₃. The precipitated solid was filtered off and washed with water to give 0.226 g of product, mp 196-198°C. (mass spectrum (ES) 483.5 (M+1)).

15

Example 61

**4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1,3- dimethyl
- 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide
hydrochloride**

To a solution of 0.610g (1.265 mmol) of 4-[(4-methoxybenzenesulfonyl)pyridin -3-ylmethylamino]-1,3-dimethyl-5-carboxylic acid, hydroxyamide in 40 ml of CH₂Cl₂-methanol (1:1) cooled to 10°C was added dropwise 1.51 ml of 1M hydrogen chloride in diethyl ether. The mixture was stirred at 10°C for 10 min. and allowed to warm to room temperature for 1 hr. The solvent was removed under vacuum and toluene (2ml) added twice and removed under vacuum after each addition. The residual solid was dried under vacuum to give 0.641 g of product as a solid, m.p. 170°-174°C.

25

Example 62

**4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-1H-
pyrazolo[3,4-b] pyridine -5-carboxylic acid, hydroxyamide**

Following the procedure of Example 49, the product of Example 47 is reacted with benzyl-(4-methoxybenzenesulfonyl)amine and sodium hydride to provide ethyl 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 124°-126°C.

35

- 44 -

Following the procedure of Example 50, the above ester is hydrolyzed to provide 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid. m.p. 108°-110°C.

Following the procedure of Example 52, the carboxylic acid is converted into the corresponding hydroxamic acid, 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide. m.p. 152°-154°C.

Example 63

10 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide**

Following the procedure of Example 53, the product of Example 47 is reacted with (4-methoxybenzenesulfonyl) (3-pyridinylmethyl) amine and sodium hydride to provide ethyl 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 89°-91°C.

Following the procedure of Example 54, the above ester is hydrolyzed to provide 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid. m.p. 136°-138°C.

Following the procedure of Example 60, the carboxylic acid is converted into the corresponding hydroxamic acid, 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide. m.p. 114°C(dec).

Example 64

25 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide, hydrochloride**

Following the procedure of Example 61, the product of Example 63 is converted into the corresponding hydrochloride salt. m.p. 161°C(dec).

Example 65

30 **Ethyl 4-chloro-1-phenyl-3-methyl-1H-pyrazolo [3,4-b]pyridine-5-carboxylate**

Following the procedure of Example 45, starting with 1-phenyl-3-methyl-5-aminopyrazole, diethyl[((1-phenyl-3-methyl-5-pyrazolyl)amino)methylene]malonate is obtained. m.p. 70°-72°C.

- 45 -

Following the procedure of Example 46, the methylene malonate is converted into ethyl 4-hydroxy-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 132°-134°C.

Following the procedure of Example 47, the hydroxy-ester is converted into the
5 chloro-ester, ethyl-4-chloro-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 108°-110°C.

Example 66

10 **4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-3-methyl-1H-pyrazolo[3,4b] pyridine-5-carboxylic acid, hydroxyamide**

Following the procedure of Example 49, the product of Example 65 is reacted with benzyl-(4-methoxybenzenesulfonyl)amine and sodium hydride to provide ethyl 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 164°-166°C.

15 Following the procedure of Example 50, the above ester is hydrolyzed to provide 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid. m.p. 246°-248°C.

Following the procedure of Example 52, the carboxylic acid is converted into the corresponding hydroxamic acid, 4-[benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide. m.p.
20 207°-210°C.

Example 67

25 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide**

Following the procedure of Example 53, the product of Example 65 is reacted with (4-methoxybenzenesulfonyl) (3-pyridinylmethyl) amine and sodium hydride to provide ethyl-4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. m.p. 148°-150°C.

30 Following the procedure of Example 54, the above ester is hydrolyzed to provide 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid. m.p. 235°-236°C.

Following the procedure of Example 60, the carboxylic acid is converted into the corresponding hydroxamic acid, 4-[(4-methoxybenzenesulfonyl)pyridin-3-

- 46 -

ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide. m.p. 192°-194°C.

Example 68

5 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide, hydrochloride**

Following the procedure of Example 61, the product of Example 67 is converted into the corresponding hydrochloride salt. m.p. 225°-226°C.

10

Example 69

4-[(4-Methoxybenzenesulfonyl)pyridin-2-ylmethyl amino] -1,3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

15 Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 70

20 **4-[(4-Methoxybenzenesulfonyl)pyridin-4-ylmethylamino] -1,3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide**

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

25

Example 71

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1-isopropyl - 1H-pyrazolo[3,4-b]pyridin -5-carboxylic acid, hydroxyamide.

30 Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 72

35 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1-benzyl-1H-pyrazolo[3,4-b] pyridin -5-carboxylic acid, hydroxyamide**

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

5

Example 73

4-[(4-Methoxybenzenesulfonyl)amino] -1-benzyl -3-methyl -1H-pyrazolo[3,4-b] pyridine -5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

10

Example 74

4-[(4-Methoxybenzenesulfonyl) 2-thienylmethylamino] -1, 3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

15

Example 75

4-[(4-Methoxybenzenesulfonyl) -3-thienylmethylamino] -1, 3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

20

Example 76

4-[(4-Methoxybenzenesulfonyl) pyridin-3-ylmethylamino] -1- (2,4-dimethoxyphenyl) - 3-methyl- 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

25

30

- 48 -

Example 77

4-[(4-Methoxybenzenesulfonyl)pyridin-3-yl)methylamino] -1-(2-methoxyphenyl)-3-methyl-1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide

5 Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

10 Example 78

4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, 15 hydroxyamides, the title compound may be prepared.

Example 79

4-{Methyl-[4-(phenoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, hydroxyamide

20 Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

25 Example 80

4-[Methyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, 30 hydroxyamides, the title compound may be prepared.

Example 81

4-[Methyl-(4-propyloxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 82

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 83

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-ethyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 84

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-tert-butyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

- 50 -

Example 85

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-methyl-3-tert-butyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide

Following procedures described in Examples 45-68 for the preparation of the
5 (substituted-4-amino)1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamides, the title compound may be prepared.

Example 86

10 **4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisothiazolo[5,4-b] pyridine-5-carboxylic Acid, Hydroxyamide**

To a stirred mixture of 0.366 g (8.4 mmol) of sodium hydride (60% in oil) in 10 ml of dry 1-methyl-2-pyrrolidinone was added (portionwise) 2.34 g (8.4 mmol) of methyl (4-methoxybenzenesulfonyl)pyridin-3-ylmethyl-amine. The mixture was stirred
15 at room temperature until gas evolution ceased and 1.80 g of (7.0 mmol) of ethyl 4-chloro-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate added. The mixture was heated at 80-90°C for 44 hours, the solvent removed under vacuum and the residue diluted with water. The mixture was extracted with ethyl acetate and the extract washed with 2N citric acid, H₂O, 1 N NaHCO₃, brine and dried (Na₂SO₄). The solution was
20 filtered through a thin pad of hydrous magnesium silicate and the pad washed with ethyl acetate. The filtrate was concentrated to dryness to give 2.39 g of ethyl 4-[(4-methoxybenzenesulfonyl)-pyridin-3-ylmethylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate as a yellow solid, m.p. 142-144°C.

Anal. for C₂₃H₂₂N₄O₅S₂

25 Calc: C, 55.4; H, 4.5; N, 11.2

Found: C, 55.5; H, 4.3; N, 11.1

Following the procedure of Example 54, a 2.25 g sample of the above ester was hydrolysed with KOH to give 0.46 g of 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid as a white solid,
30 m.p. 234-236°C.

Anal. for C₂₁H₁₈N₄O₅S₂

Calc: C, 53.6; H, 3.9; N, 11.9

Found: C, 53.5; H, 3.8; N, 11.8

35

- 51 -

Following the procedure of Example 60, 2.0 g of the preceding compound as the potassium salt was converted to the title compound to give 0.39 g of off-white solid, m.p. 145-149°C. The hydrochloride salt was prepared from 0.27 g of the hydroxyamide according to the procedure of Example 61 to give 0.26 g of yellow solid, m.p. 224°C dec.

Anal. for $C_{21}H_{19}N_5O_5S_2 \cdot HCl$

Calc: C, 48.3; H, 3.9; N, 13.4

Found: C, 48.0; H, 3.8; N, 13.2

Example 87

4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic Acid, Hydroxyamide

Following the procedure of Example 86, 1.7 g (7 mmol) of ethyl 4-chloro-3-methylisoxazolo[5,4-b]pyridine-5-carboxylate was reacted with 2.92 g (0.0105 mmol) of (4-methoxybenzenesulfonyl)pyridin-3-ylmethylamine to give 1.01 g of ethyl 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisoxazolo[5,4-b]pyridine-5-carboxylate as a white solid, m.p. 128-130°C.

Anal. for $C_{23}H_{22}N_4O_6S$:

Calc: C, 57.3; H, 4.6; N, 11.6

Found: C, 57.3; H, 4.7; N, 11.5

A mixture of 1.01 g (2.1 mmol) of the preceding compound in 10 ml of tetrahydrofuran and 2.93 ml of 1 N NaOH was stirred at room temperature overnight and the solvent removed. The residue was diluted with H_2O and acidified with 2N citric acid (pH 4). The solid was filtered off and washed with H_2O to give 0.88 g of 4-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic acid as a white solid, m.p. 244-246°C.

Anal. for $C_{21}H_{18}N_4O_6S$:

Calc: C, 55.5; H, 4.0; N, 12.3

Found: C, 55.2; H, 4.0; N, 12.2

Following the procedure of Example 60, a 0.86 g (1.89 mmol) sample of the preceding compound was converted to the title compound to give 0.42 g of off-white solid, m.p. 150°C dec.

Anal. for $C_{21}H_{19}N_5O_6S$:

- 52 -

Calc: C, 53.7; H,4.1; N,14.9
Found: C, 53.4; H,4.5; N,14.4

The hydrochloride salt was prepared according to the procedure of Example 61
5 from 0.25 g of the title compound to give a solid which on trituration with ethyl acetate
gave 0.27 g of off-white solid, m.p. 212-215°C.

Anal. for $C_{21}H_{19}N_5O_6S \cdot HCl$:

Calc: C, 49.8; H,4.0; N,13.8
Found: C, 49.4; H,4.1; N,14.0

10

Example 88

7-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic Acid, Hydroxyamide

15 Following the procedure of Example 86, 1.8 (7.5 mmol) of ethyl 7-chloro-2-methyl pyrazolo[1,5-a] pyrimidine-6-carboxylate was reacted with 2.92 g (10.5 mmol) of (4-methoxybenzenesulfonyl)pyridin-3-ylmethyl-amine to give 1.64 g of ethyl 7-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid as a yellow solid, m.p. 108-110°C.

20 Anal. for $C_{23}H_{23}N_5O_5S$

Calc: C, 57.4; H,4.8; N,14.5
Found: C, 54.5; H,4.7; N,14.4

A mixture of 1.54 g (3.20 mmol) of the preceding compound, tetrahydrofuran
25 (15 ml) and 4.15 ml of 1 N NaOH was stirred at room temperature overnight and the solvent removed under vacuum. The residue was diluted with H_2O and extracted with diethyl ether and ethyl acetate. The aqueous layer was acidified with 2 N citric acid (pH 5) and the solid filtered off and washed with H_2O . The solid was dried at 76°C in a vacuum oven to give 1.03 g of 7-[(4-methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid as an off-white
30 solid, m.p. 249-251°C.

Anal. for $C_{21}H_{19}N_5O_5S$

Calc: C, 55.6; H,4.2; N,15.4
Found: C, 55.2; H,4.2; N,15.6

35

- 53 -

A mixture of 1.0 g (2.2 mmol) of the preceding compound, 3 ml of $\text{CH}_3\text{OH}\cdot\text{H}_2\text{O}$ (2:1) and 2.43 ml of 1 N KOH was stirred 0.5 hours and the solvent removed under vacuum. Toluene (10 ml) was added three times and the solvent removed after each addition. The residue was dried in a vacuum oven and following the procedure of Example 60 the potassium salt was converted to the title compound to obtain 0.29 g of yellow solid, m.p. 185°C dec.

Anal. for $\text{C}_{21}\text{H}_{20}\text{N}_6\text{O}_5\text{S}$

Calc: C, 53.8; H, 4.3; N, 17.9

Found: C, 53.9; H, 4.4; N, 17.3

10

The hydrochloride salt was prepared according to the procedure of Example 61 from 0.18 g of the title compound to give 0.22 g of yellow solid, m.p. 170°C dec.

Anal. for $\text{C}_{21}\text{H}_{20}\text{N}_6\text{O}_5\text{S}\cdot\text{HCl}$

15 Calc: C, 50.0; H, 4.2; N, 16.6

Found: C, 48.7; H, 4.4; N, 16.1

Example 89

20 4-[[4-(4-Chlorophenyloxy)benzenesulfonyl] methylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic Acid, Hydroxyamide

To 20 ml of a 2 molar solution of methylamine in tetrahydrofuran chilled in an ice bath was added a solution of 5.16 g of 4-(4-chlorophenyloxy)benzene-sulfonyl chloride in 50 ml of tetrahydrofuran and 20 ml of CH_2Cl_2 . The mixture was refluxed for 2 hours and the solvent removed under vacuum and the residue partitioned between 25 $\text{CH}_2\text{Cl}_2\cdot\text{H}_2\text{O}$ (1:1). The CH_2Cl_2 layer was separated and washed with 2 N citric acid, brine and dried (Na_2SO_4). The solution was filtered through a thin pad of hydrous magnesium silicate and the filter pad washed with CH_2Cl_2 . The filtrate was concentrated to give 4.5 g of N-methyl-4-(4-chlorophenoxy)benzene-sulfonamide as a yellow solid, m.p. 80-83°C.

30 Following the procedure of Example 53, 1.79 g (6 mmol) of the preceding compound was reacted with 1.29 g (5 mmol) of ethyl 4-chloro-1,3-dimethyl pyrazolo[3,4-b]pyridine-5-carboxylate at 80-90°C for 48 hours to give 3.12 g of solid. Chromatography on silica gel with hexane-diethyl ether (2:1) as eluent to give 1.92 g of ethyl 4-[[4-(4-chlorophenyloxy)benzenesulfonyl]methylamino]-1,3-dimethyl-1H-35 pyrazolo[3,4-b]pyridine-5-carboxylate as an off-white solid, m.p. 52-55°C.

- 54 -

Anal. for $C_{24}H_{23}ClN_4O_5S$

Calc: C, 56.0; H, 4.5; N, 10.9

Found: C, 55.9; H, 4.4; N, 10.6

- 5 A mixture of 1.90 g (3.69 mmol) of the preceding ester and 4.05 ml (4.06 mmol) of 1 N KOH in 15 ml of tetrahydrofuran was stirred at room temperature for 2 days and then 0.75 ml of 1 N KOH was added and the mixture was refluxed overnight. The solvent was removed under vacuum and the toluene (25 ml) added to the residue and removed under vacuum. The residue was triturated with ethyl acetate and the solid
10 filtered and dried under vacuum to give 1.6 g of 4-[[4-(4-chlorophenoxy)benzenesulfonyl]methylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, potassium salt. Following the procedure of Example 52, 1.60 g of the preceding potassium salt was converted to the title compound to obtain 1.82 g of solid. This solid was purified on thick layer silica gel plates with 4% CH_3OH in ethyl acetate as solvent to
15 give 0.59 g of light yellow solid, m.p. 120°C dec.

Anal. for $C_{22}H_{20}ClN_5O_5S \cdot H_2O$

Calc: C, 51.7; H, 4.3; N, 13.7

Found: C, 51.2; H, 4.0; N, 13.6

20

Example 90

4-[[4-(4-Chlorophenoxy)benzenesulfonyl] methylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic Acid Hydroxyamide

- Following the procedure of Example 86, 1.17 g (38.5 mmol) of N-methyl-4-(4-chlorophenoxy)benzene-sulfonamide was reacted with 0.916 g (3.57 mmol) of ethyl 4-chloro-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate for 33 hours to give a solid which was triturated with ethyl acetate to give 0.99 g of ethyl 4-[[4-(4-chlorophenoxy)benzenesulfonyl]methylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate as a white solid, m.p. 117-120°C.

- 30 Anal. for $C_{23}H_{20}ClN_3O_5S_2$

Calc: C, 53.3; H, 3.9; N, 8.1

Found: C, 53.2; H, 3.9; N, 7.8

- Following the procedure of Example 54, a mixture of 0.96 g (1.85 mmol) of the
35 preceding compound and 2.40 ml of 1 N KOH in 15 ml of tetrahydrofuran-ethanol (2:1)

- 55 -

- was refluxed 16 hours and the solvent removed under vacuum. Toluene (25 ml) was added to the residue and the solvent removed. The residue was triturated with ethyl acetate to give a solid which was dried under vacuum to give 0.80 g of 4-[[4-(4-chlorophenoxy)-benzenesulfonyl] methylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid, potassium salt. Following the procedure of Example 52, 0.80 g (1.51 mmol) of the preceding compound was converted to the title compound to give 0.82 g of solid. This solid was chromatographed on thick layer silica gel plates with 4% CH₃OH in ethyl acetate as solvent to give 0.19 g of solid which was triturated with hexane to give 0.12 g of 115°C dec.
- Anal. for C₂₁H₁₇ClN₄O₅S₂•1/4 hexane
- Calc: C, 51.3; H, 3.9; N, 10.6
- Found: C, 51.9; H, 4.2; N, 10.4

Example 91

4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic Acid, Hydroxyamide

- A suspension of 35 g of [4-(4-pyridinyloxy) benzenesulfonyl chloride hydrochloride in 800 ml of tetrahydrofuran was added slowly to 163 ml of 2 M methylamine in tetrahydrofuran. An additional 550 ml of tetrahydrofuran was added and the suspension refluxed for 3 hours. The solvent was removed under vacuum and the residue partitioned between CH₂Cl₂ and H₂O. The organic layer was separated, washed with saturated NaHCO₃, brine and dried (Na₂SO₄). The solvent was removed to give 13.11 g of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide, m.p. 125-127°C. The aqueous layer was neutralized with 1 N NaOH (solid precipitated) and extracted with CH₂Cl₂. The extract was washed with brine and dried (Na₂SO₄). The solvent was removed to give an additional 15.9 g of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide. A 2.80 g (10.6 mmol) portion of the preceding compound was added to a stirred suspension of 0.424 g (10.6 mmol) of sodium hydride (60% in oil) in 20 ml of dry 1-methylpyrrolidinone. The mixture was stirred until gas evolution ceased and 2.8 g (10.0 mmol) of ethyl 4-chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate was added followed by the addition of 20 ml of dry 1-methylpyrrolidinone. The mixture was heated at 80-90°C for 58 hours. The solvent was removed under vacuum and the residue extracted with ethyl acetate (100 ml) and the extract washed with 100 ml of H₂O. The extract was stirred with 40 ml of 1 N HCl for 1 hour and the aqueous layer separated and neutralized (pH 6-7) with 1 N NaHCO₃.

- 56 -

The mixture was extracted with ethyl acetate and the extract washed with H₂O, brine and dried (Na₂SO₄). The solution was filtered through a thin pad of hydrous magnesium silicate and the filter pad washed with ethyl acetate. The filtrate was concentrated to dryness and the residue crystallized with ethyl acetate-hexane to give 3.43 g of ethyl 4-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate as pale yellow crystals, m.p. 117-119°C.

Anal. for C₂₃H₂₃N₅O₅S

Calc: C, 57.4; H, 4.8; N, 14.5

Found: C, 57.3; H, 4.7; N, 14.5

10

A mixture of 7.44 g (15.5 mmol) of the preceding compound and 18.54 ml of 1 N KOH in 100 ml tetrahydrofuran-ethanol (6:4) was refluxed overnight under nitrogen. The solvent was removed and toluene and ethanol added and the solvent removed under vacuum. The residue was triturated with diethyl ether, filtered and the solid dried under vacuum to give 7.57 g of the potassium salt of 4-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid. A 0.18 g sample of the preceding compound was dissolved in H₂O and the solution brought to pH 6 with 2 N citric acid. The precipitated solid was filtered, washed with H₂O and dried under vacuum to give 0.045 g of 4-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid as a white solid, m.p. 171-185°C.

15

20

Anal. for C₂₁H₁₉N₅O₅S•2 H₂O

Calc: C, 51.5; H, 4.7; N, 14.3

Found: C, 51.8; H, 4.1; N, 14.2

25

To a chilled (0°C) mixture of 2.2 ml (4.4 mmol) of oxalyl chloride in CH₂Cl₂ (2 M) was added dropwise 0.308 ml (4.0 mmol) of N,N-dimethylformamide followed by the addition of 0.982 g (2 mmol) of the preceding potassium salt and 10 ml of CH₂Cl₂ (Mixture A). A mixture of 0.444 ml (8 mmol) of hydroxylamine in H₂O (50% w), 0.634 ml of triethylamine and 3.50 ml of tetrahydrofuran was chilled and added to the chilled Mixture A. After stirring at 0°C for 15 min., the mixture was stirred at room temperature overnight. The solvent was removed and the residue partitioned between 10 ml of CH₂Cl₂ and 30 ml of H₂O. The solid was filtered off and suspended in H₂O. The pH of the mixture was adjusted to pH 6-7 with 2 N citric acid, stirred, filtered and the solid washed with H₂O and CH₂Cl₂ to give 0.47 g of 4-{methyl-[4-(4-

30

35

- 57 -

pyridinyloxy)benzenesulfonyl]amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide as a white solid, m.p. 147° dec.

Anal. for $C_{21}H_{20}N_6O_5S \cdot H_2O$

Calc: C, 51.8; H, 4.6; N, 17.3

5 Found: C, 51.3; H, 4.8; N, 17.4

To a cooled 2.2 g (4.7 mmol) sample of the preceding compound in 40 ml of CH_2Cl_2 - CH_3OH (1:1) was added 5.64 ml of 1 M HCl in diethyl ether. The mixture was stirred in an ice bath for 10 min. and then at room temperature for 1 hour. The solvent was removed under vacuum and 25 ml of toluene added (two times) and removed under vacuum. The solid was triturated with 20 ml of methanol to give 1.60 g of the product of the Example as a white solid, m.p. 197-200°C.

Anal. for $C_{21}H_{20}N_6O_5S \cdot HCl \cdot H_2O$

Calc: C, 48.2; H, 4.4; N, 16.1

15 Found: C, 47.7; H, 4.7; N, 15.9

Example 92

4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic Acid, Hydroxyamide

20 Following the procedure of Example 86, 1.28 g (5 mmol) of ethyl 4-chloro-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate was reacted with 1.4 g (5.3 mmol) of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide to give 2.78 g of an oil. This oil was dissolved in 10 ml of ethyl acetate and 13 ml of 1 N HCl added. The organic layer was separated. The aqueous layer was brought to pH 6 with 1 N NaOH and extracted with 25 40 ml of CH_2Cl_2 . The CH_2Cl_2 extract was washed with brine and dried (Na_2SO_4). The solvent was removed to give 1.69 g of solid which was chromatographed on silica gel with ethyl acetate as solvent to give 0.95 g of ethyl 4-{methyl-[4-(4-pyridinyloxy)benzene-sulfonyl]amino}-3-methylisothiazolo[5,4-b]pyridine-5-carboxylate as a yellow solid, m.p. 57-60°C.

30 Anal. for $C_{22}H_{20}N_4O_5S_2$

Calc: C, 54.5; H, 4.2; N, 11.6

Found: C, 54.4; H, 4.1; N, 11.2

Following the procedure of Example 50, a mixture of 0.95 g of the preceding ester and 2.8 ml of 1 N KOH in 2.5 ml of tetrahydrofuran and 25 ml of ethanol was 35

- 58 -

refluxed 24 hours and the solvent removed under vacuum to give 1.0 g of solid. Following the procedure of Example 91, a 0.82 g sample of the resulting potassium salt was reacted with hydroxylamine to give 0.78 g of solid. Chromatography on thick layer silica gel plates with ethyl acetate-CH₃OH (85:15) as solvent gave 0.22 g of 4-{methyl-

- 5 [4-(4-pyridinyloxy)benzenesulfonyl]amino}-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide as an off-white solid, m.p. 146°C dec.

Anal. for C₂₀H₁₇N₅O₅S₂•1/2 H₂O

Calc: C, 50.0; H, 3.8; N, 14.6

Found: C, 49.7; H, 4.0; N, 13.8

10

To a solution of 0.20 g of the preceding compound in 3.4 ml of CH₂Cl₂-CH₃OH (1:1) was added 0.508 ml of 1 N HCl in diethyl ether. The solvent was removed under vacuum and the solid dried 20 hours under vacuum to give the hydrochloride salt of the title compound as an off-white solid, m.p. 186°C dec.

- 15 Anal. for C₂₀H₁₇N₅O₅S₂•HCl•H₂O

Calc: C, 45.6; H, 3.8; N, 13.3

Found: C, 45.6; H, 4.1; N, 12.7

Example 93

- 20 **4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-1-methyl-3-phenyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, Hydroxyamide**

Following the procedure described in Example 91, 1.45 g (5.5 mmol) of N-methyl-4-(pyridinyl-4-oxy)benzenesulfonamide was reacted with 1.58 g (5 mmol) of ethyl 4-chloro-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate in 20 ml of

25 1-methyl-pyrrolidinone to give 1.62 g of ethyl 4-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]-amino}-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate as an off-white solid, m.p. 128-131°C.

Anal. for C₂₈H₂₅N₅O₅S

Calc: C, 61.9; H, 4.6; N, 12.9

- 30 Found: C, 61.7; H, 4.7; N, 12.8

A mixture of 1.60 g (2.94 mmol) of the preceding compound and 3.53 ml of 1 N KOH in 8 ml of tetrahydrofuran-ethanol (1:1) was refluxed overnight and the solvent removed. The residue was triturated with diethyl ether to give 1.62 g of potassium 4-

- 59 -

{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylate as solid.

5 Following the procedure of Example 91, a 1.47 g (2.66 mmol) sample of the preceding compound was reacted with hydroxylamine to give 0.49 g of solid. This solid (0.45 g) was dissolved in 15 ml of CH₂Cl₂-CH₃OH (1:1) and to the chilled solution was added 2.26 ml of 1 N HCl in diethyl ether. The mixture was stirred at room temperature for 1 hour and the solvent removed to give the hydrochloride salt of the title compound as an off-white solid, m.p. 195°C dec.

10 Anal. for C₂₆H₂₂N₆O₅S•HCl•H₂O

Calc: C, 53.4; H, 5.0; N, 14.4

Found: C, 53.2; H, 4.3; N, 14.3

Example 94

15 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic acid, Hydroxyamide,

20 Following the procedure of Example 91, 1.45 g (5.5 mmol) of N-methyl-4-(pyridinyl-4-oxy)benzenesulfonamide was reacted with 1.20 g (5 mmol) of ethyl 4-chloro-3-methylisoxazolo[5,4-b]pyridine-5-carboxylate to give 1.29 g of ethyl 4-{methyl-[4-(4-pyridinyloxy)-benzenesulfonyl]amino}-3-methylisoxazolo[5,4-b]pyridine-5-carboxylate as an off-white solid, m.p. 155-157°C.

Anal. for C₂₂H₂₀N₄O₆S•1/2 H₂O

Calc: C, 55.3; H, 4.4; N, 11.7

Found: C, 55.2; H, 4.3; N, 11.6

25

30 A mixture of 1.84 g (3.92 mmol) of the preceding ester and 4.71 ml of 1 N KOH in 10 ml of tetrahydrofuran-ethanol (1:1) was refluxed overnight and the solvent removed under vacuum. The residue was triturated with diethyl ether and filtered to give 1.59 g of potassium 4-{methyl-[4-(4-pyridinyloxy)benzene-sulfonyl]amino}-3-methylisoxazolo[5,4-b]pyridine-5-carboxylate as a pale yellow solid. Following the procedure of Example 91, the preceding potassium salt (1.45 g) was reacted with hydroxylamine to give a solid. The solid was partitioned between 30 ml of CH₂Cl₂ and 30 ml of H₂O and filtered. The CH₂Cl₂ layer was stirred with 2 N citric acid (aqueous phase-pH 6). The solid which precipitated was filtered off and suspended in H₂O. The
35 aqueous suspension was adjusted to pH 7 with 1 N NaHCO₃ and filtered. The solid

- 60 -

was again suspended in H₂O and the pH of the aqueous suspension adjusted to pH 6 with 2 N citric acid. The mixture was stirred and filtered to give 0.66 g of 4-{methyl-[4-(4-pyridinyl-oxy)benzenesulfonyl]amino}-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide, m.p. 141°C dec.

5 Anal. for C₂₆H₂₂N₆O₅S•H₂O

 Calc: C, 58.9; H,4.4; N,15.3

 Found: C, 56.3; H,4.3; N,15.2

10 To a chilled solution of 0.569 g (1.25 mmol) of the preceding hydroxyamide in 10 ml of CH₂Cl₂-CH₃OH (1:1) was added 1.50 ml of 1 N HCl in diethyl ether. The mixture was stirred at room temperature for 1 hour and the solvent removed to give 0.582 g of the hydrochloride salt of the title compound as a light brown solid, m.p. 170°C dec.

 Anal. for C₂₀H₁₇N₅O₆S•HCl•H₂O

15 Calc: C, 47.1; H,4.0; N,13.7

 Found: C, 47.2; H,4.5; N,13.2

Example 95

20 **7-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl] amino}-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid, hydroxyamide**

 Following the procedure of Example 86, 1.19 g (5 mmol) of ethyl 7-chloro-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylate was reacted with 1.45 g (5.5 mmol) of N-methyl-4-(pyridinyl-4-oxy)benzenesulfonamide (as the sodium salt) in 20 ml of 1-methylpyrrolidinone at 80-90°C for 66 hours. The solvent was removed and the residue
25 partitioned between 30 ml of ethyl acetate and 30 ml of H₂O. The organic layer was separated and stirred with 30 ml of 1 N HCl for 1 hour and filtered. The aqueous layer was separated and the pH adjusted to pH 6 with saturated NaHCO₃ solution. The mixture was extracted with ethyl acetate and the extract washed with brine and dried (Na₂SO₄). The solution was filtered through a thin pad of hydrous magnesium silicate
30 and the filter pad washed with ethyl acetate. The filtrate was concentrated to dryness and the solid suspended in 50 ml of H₂O and the pH of the stirred suspension adjusted to pH 7 with 1 N NaHCO₃. The mixture was extracted with ethyl acetate and the extract washed with brine and dried (Na₂SO₄). The solution was filtered through a thin pad of hydrous magnesium silicate and the filtrate concentrated to dryness to give 1.35 g of

- 61 -

ethyl 7-[methyl-[4-(4-pyridinyloxy)-benzenesulfonyl]amino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylate as a yellow solid, m.p. 139-144°C.

Anal. for $C_{22}H_{21}N_4O_5S$

Calc: C, 56.5; H, 4.5; N, 15.0

5 Found: C, 56.8; H, 4.6; N, 14.9

A mixture of 1.25 g (2.67 mmol) of the preceding ester and 3.21 ml of 1 N KOH in 15 ml of tetrahydrofuran was stirred at room temperature overnight. The mixture was filtered and the solid washed with diethyl ether and ethyl acetate to give 0.5 g of potassium 7-[methyl-[4-(4-pyridinyloxy)benzene-sulfonyl]amino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylate. A 0.15 g sample of the preceding compound was dissolved in water and the pH adjusted to pH 6 with 2 N citric acid. The mixture was filtered to give 0.12 g of 7-[methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid as a white solid, m.p. 246-248°C.

Anal. for $C_{20}H_{17}N_5O_5S$

Calc: C, 54.7; H, 3.9; N, 15.9

Found: C, 54.2; H, 4.1; N, 16.4

20 Following the procedures of Example 91, the title compound is prepared from the carboxylic acid described above.

Example 96

25 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-7-methyl-1,8-naphthyridine-3-carboxylic Acid, Hydroxyamide

A mixture of 10.8 g (0.1 M) of 2-amino-6-methylpyridine and 23.78 g (0.12 M) of diethyl ethoxymethylenemalonate was heated in an oil bath (preheated to 90°C) for 1 hr. The mixture was cooled to room temperature and the solid was recrystallized from 100 ml of ethanol to give 26.38 g of 2-(2,2-dicarbethoxy-1-vinylamino)-6-methylpyridine, m.p. 102-104°C. Reported m.p. 107-108°C (U.S. Patent 4,166,817 issued Sept. 4, 1979). A solution of 20.79 g of the preceding compound in 40 ml of Dowtherm (heated to 80°C) was dropped into 100 ml of Dowtherm which had been preheated to 258°C. The temperature of the mixture dropped to 200°C and after the temperature returned to 250°C (10 min), the mixture was heated at 250°C for 30 min.

35 The mixture was immediately cooled to room temperature and allowed to stand

- 62 -

overnight. The mixture was filtered and the solid washed with hexane and CH_2Cl_2 to give 4.24 g of ethyl 4-hydroxy-7-methyl-1,8-naphthyridine-3-carboxylate as a brown solid.

A mixture of 4.2 g (0.018 M) of the preceding compound and 45 ml of POCl_3 (0.48 M) was heated at 70-80°C for 4 hours. The solution was concentrated under vacuum and the residue poured onto crushed ice. The resulting mixture was neutralized with 5 N NaOH to pH 6 and extracted with diethyl ether (3x250 ml). The extract was washed with brine and dried over Na_2SO_4 . The solvent was removed under vacuum to give 2.24 g of ethyl 4-chloro-7-methyl-1,8-naphthyridine-3-carboxylate as a yellow solid, m.p. 72-74°C: reported m.p. 92-93°C (dec) (U.S. Patent 4,166,817).

To a mixture (under nitrogen) of 0.22 g (5.5 mmol) of NaH (60% in oil) in 20 ml of 1-methylpyrrolidinone was added 1.45 g (5.5 mmol) of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide. After gas evolution ceased, 1.25 g (5 mmol) of ethyl 4-chloro-7-methyl-1,8-naphthyridine-3-carboxylate was added. The mixture was heated at 80-90°C for 60 hrs and the solvent removed under vacuum. The residue was partitioned between H_2O and ethyl acetate and the mixture filtered through diatomaceous earth. The organic layer of the filtrate was separated and washed with H_2O and dried (Na_2SO_4). The solution was filtered through a thin pad of hydrous magnesium silicate and the filtrate concentrated under vacuum to give 0.97 g of a brown solid (mass spectrum 265 (75%); 479 (25%). The product (mass spectrum (M+H) 479) is separated by chromatography on silica gel and following the procedure of Example 91 is converted to the title compound.

Example 97

25 7-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-2,3-dimethylimidazo[4,5-b]pyridine-6-carboxylic Acid, Hydroxyamide

Following the general procedure described in J. Chem. Soc. Perkin Trans. 1, 2789 (1992) a mixture of 1,2-dimethyl-5-nitroimidazole (8.46 g; 0.06 M), diethyl ethoxymethylenemalonate (13.08 g; 0.06 M) and 2.11 g of 5% Pd on carbon in 135 ml of dioxane was reduced in a Parr Hydrogenator at 35 to 40 psi of hydrogen for 29 hours. The mixture was filtered through diatomaceous earth and the solvent removed to give a brown oil. This oil was dissolved in 100 ml of 2 N HCl and the pH adjusted to pH 5 with 10 N NaOH. The mixture was extracted twice with 100 ml of ethyl acetate (extract discarded). The pH was adjusted to pH 7 and extracted with 150 ml of ethyl acetate and then the pH was adjusted to pH 9 and again extracted twice with 150 ml of

- 63 -

ethyl acetate. The pH 7 and pH 9 extracts were combined and washed with brine and dried over Na_2SO_4 . The solution was filtered through a thin pad of hydrous magnesium silicate and the filtrate concentrated to dryness to give 7.61 g of 5-[2,2-bis(ethoxycarbonyl)-1-vinylamino]-1,2-dimethylimidazole (diethyl [(1,2-dimethylimidazol-5-yl)aminomethylene]malonate) as a brown oil.

A mixture of the preceding compound (7.9 g) and 35 ml of POCl_3 was refluxed for 7 hours under nitrogen and then concentrated under vacuum. The black residue was poured onto crushed ice (with stirring) and the mixture brought to pH 5 with 5 N NaOH. The mixture was extracted with 150 ml of ethyl acetate, 200 ml of diethyl ether and 200 ml of CH_2Cl_2 . Each extract was washed with 1 N NaHCO_3 , brine and dried (Na_2SO_4). The solutions were combined and filtered through a thin pad of hydrous magnesium silicate. The filtrate was concentrated to dryness under vacuum to give 4.1 g of ethyl 7-chloro-2,3-dimethylimidazo[4,5-b]pyridine-6-carboxylate as a tan solid, m.p. 85-90°C. A sample crystallized from diethyl ether gave crystals, m.p. 117-119°C, Anal. for $\text{C}_{11}\text{H}_{12}\text{ClN}_3\text{O}_2 \cdot 1/2 \text{H}_2\text{O}$
Calc. C, 48.8; H, 4.6; N, 15.9
Found C, 50.3; H, 5.6; N, 16.0

Following the procedure of Example 91, a 0.759 g (3 mmol) portion of the preceding ester was heated at 80-90°C in 14 ml of 1-methylpyrrolidinone with the sodium salt of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide [from 0.871 g; 3.3 mmol of N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide and 79.2 mg (3.3 mmol) of NaH (60% in oil)]. The mixture was heated for 2.5 days at 80-90°C and 3 hrs at 100°C and then the solvent was removed under vacuum. The residue was extracted with ethyl acetate and the extract washed with H_2O . The organic layer was stirred with 1 N HCl (30 ml) for 0.5 hour and the aqueous layer separated and brought to pH 6 with 5 N NaOH. The mixture was extracted with ethyl acetate and the extract dried over Na_2SO_4 . The solvent was removed under vacuum to give 1.0 g of a brown oil which contained the product ethyl 7-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-2,3-dimethylimidazo[4,5-b]pyridine-6-carboxylate; mass spectrum (ES) 479.1 (M+H). Following the procedure of Example 91, the preceding compound is converted to the title compound.

- 64 -

Example 98

**2-Methyl-4-{methyl-[4-(4-pyridinyloxy)
benzenesulfonyl]amino}thieno[3,4-b]pyridine-3-carboxylic Acid,
Hydroxyamide**

5 A mixture of 10.0 g (63.6 mmol) of methyl 3-aminothiophene-4-carboxylate, 10.1 g (63.6 mmol) of ethyl (trans)-3-ethoxycrotonate and 40 mg of p-toluenesulfonic acid, monohydrate in 50 ml of p-xylenes was refluxed overnight and the solvent removed under vacuum. To the residue was added 20 ml of p-xylenes, and 23.7 ml of NaOC₂H₅ (21% by wt) (63.6 mmol) in ethanol and the mixture refluxed for 3 hrs. The
10 solvent was removed, the residue diluted with H₂O and the pH adjusted to pH 4 with 1 N HCl. The precipitate was filtered, washed with water and ethyl acetate to give 4.95 g of 4-hydroxy-2-methyl-thieno[3,4-b]pyridine-3-carboxylic acid as a brown solid.

 The preceding compound (1.4 g) was dissolved in 10 ml of dry methanol and HCl gas bubbled into the solution for 10 min. The solution was stirred overnight at
15 room temperature and the solvent removed under vacuum. The residue was dissolved in ethyl acetate and the solution washed with saturated NaHCO₃, brine and dried (Na₂SO₄). The solvent was removed to give a solid which was triturated with ethyl acetate. The mixture was cooled and filtered to give 0.765 g of methyl 4-chloro-2-methylthieno[3,4-b]pyridine-3-carboxylate as a yellow solid.

20 Following the procedure of Example 91, the preceding compound is converted to the title compound.

Example 99

25 **5-Methyl-7-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-
thieno[3,2-b]pyridine-6-carboxylic Acid, Hydroxyamide**

 Following the procedure described in J. Med. Chem. 33, 2640 (1990), a mixture of 10 g (63.6 mmol) of methyl 3-aminothiophene-2-carboxylate (10.1 g) (63.6 mmol) of ethyl (trans)-3-ethoxycrotonate and 40 mg of p-toluenesulfonic acid monohydrate in 80 ml of xylene was refluxed overnight. The solvent was removed
30 under vacuum and the residue dissolved in ethyl acetate. The solution was washed with H₂O, 2 N citric acid, 1 N NaHCO₃, brine and dried (Na₂SO₄). The solid 16 g was chromatographed on silica gel with hexane-ethyl acetate (5:1) to give 6.65 g of ethyl 3-[(2-methoxycarbonyl-3-thienyl)amino]crotonate as a yellow oil. To a sample of 0.269 g (1 mmol) of the preceding compound in 3.5 ml of xylenes (chilled in an ice bath) was
35 added 44 mg (1.1 mmol) of NaH (60% in oil). The mixture was refluxed for 3 hours

and the solvent removed. The residue diluted with water and extracted with ethyl acetate. The aqueous layer was acidified (1 N HCl) to pH 4 and the mixture extracted with ethyl acetate. The extract was washed with brine, dried over Na₂SO₄ and the solvent removed to give 190 mg of a mixture (1:1) of methyl 7-hydroxy-5-methylthieno[3,2-b]pyridine-6-carboxylate and ethyl 7-hydroxy-5-methylthieno[3,2-b]pyridine-6-carboxylate as a solid. The preceding ethyl ester was prepared in the following manner.

A mixture of 5.0 g (31.8 mmol) of methyl 3-aminothiophene-2-carboxylate, 5.03 g (31.8 mmol) of ethyl (trans)-3-ethoxycrotonate and 20 mg of p-toluenesulfonic acid monohydrate in 50 ml of p-xylenes was refluxed 1 hour and allowed to stand 2 days at room temperature. The mixture was concentrated under vacuum and then cooled (ice bath). To the solution was added 12.4 ml of a solution of sodium ethoxide (21% by wt) in ethanol. The mixture was refluxed for 2 hours and the solvent removed. The residue was partitioned between H₂O and diethyl ether and the H₂O layer separated and acidified to pH 4 with 1 N HCl. The mixture was extracted with ethyl acetate and the extract washed with brine and dried (Na₂SO₄). The solvent was removed to give 2.2 g of brown solid. The solid was triturated with ethyl acetate, chilled and filtered to give 1.0 g of ethyl 7-hydroxy-5-methylthieno[3,2-b]pyridine-6-carboxylate as a light tan solid (mass spectrum (ES) 238 (M+H)).

A mixture of the preceding compound (0.985g) and 4 ml of POCl₃ was refluxed 2 hours and the mixture poured onto crushed ice. The mixture was extracted with ethyl acetate and the extract concentrated to dryness. The residue was dissolved in CH₂Cl₂ and the solution washed with H₂O and dried over Na₂SO₄. The solution was filtered through a thin pad of hydrous magnesium silicate and the filtrate concentrated to dryness to give 0.62 g of ethyl 7-chloro-5-methylthieno[3,2-b]pyridine-6-carboxylate as a yellow oil; thin layer chromatography on silica gel; R_f = 0.9; ethyl acetate-hexane (1:1).

Following the procedure of Example 86, the preceding compound is reacted with N-methyl-4-(pyridin-4-yloxy)benzenesulfonamide to give ethyl 5-methyl-7-{methyl-[4-(4-pyridinyloxy) benzenesulfonyl]amino}thieno[3,2-b]pyridine-6-carboxylate. As described for Example 91, the preceding compound is converted to the title compound.

Representative compounds of this invention were evaluated as inhibitors of the enzymes MMP-1, MMP-9, MMP-13 and TNF- α converting enzyme (TACE). The standard pharmacological test procedures used, and results obtained which establish this biological profile are shown below.

- 66 -

Test Procedures for Measuring MMP-1, MMP-9, and MMP-13 Inhibition

These standard pharmacological test procedures are based on the cleavage of a thiopeptide substrates such as Ac-Pro-Leu-Gly(2-mercapto-4-methyl-pentanoyl)-Leu-Gly-OEt by the matrix metalloproteinases MMP-1, MMP-13 (collagenases) or MMP-9 (gelatinase), which results in the release of a substrate product that reacts colorimetrically with DTNB (5,5'-dithiobis(2-nitro-benzoic acid)). The enzyme activity is measured by the rate of the color increase. The thiopeptide substrate is made up fresh as a 20 mM stock in 100% DMSO and the DTNB is dissolved in 100% DMSO as a 100 mM stock and stored in the dark at room temperature. Both the substrate and DTNB are diluted together to 1 mM with substrate buffer (50 mM HEPES pH 7.5, 5 mM CaCl₂) before use. The stock of enzyme is diluted with buffer (50 mM HEPES, pH 7.5, 5 mM CaCl₂, 0.02% Brij) to the desired final concentration. The buffer, enzyme, vehicle or inhibitor, and DTNB/substrate are added in this order to a 96 well plate (total reaction volume of 200 μ l) and the increase in color is monitored spectrophotometrically for 5 minutes at 405 nm on a plate reader and the increase in color over time is plotted as a linear line.

Alternatively, a fluorescent peptide substrate is used. In this test procedure, the peptide substrate contains a fluorescent group and a quenching group. Upon cleavage of the substrate by an MMP, the fluorescence that is generated is quantitated on the fluorescence plate reader. The assay is run in HCBC assay buffer (50mM HEPES, pH 7.0, 5 mM Ca⁺², 0.02% Brij, 0.5% Cysteine), with human recombinant MMP-1, MMP-9, or MMP-13. The substrate is dissolved in methanol and stored frozen in 1 mM aliquots. For the assay, substrate and enzymes are diluted in HCBC buffer to the desired concentrations. Compounds are added to the 96 well plate containing enzyme and the reaction is started by the addition of substrate. The reaction is read (excitation 340 nm, emission 444 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line.

For either the thiopeptide or fluorescent peptide test procedures, the slope of the line is calculated and represents the reaction rate. The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($\bar{x} \pm \text{sem}$) of the control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple

- 67 -

comparison test. Dose-response relationships can be generated using multiple doses of drug and IC₅₀ values with 95% CI are estimated using linear regression.

In vivo MMP Inhibition Test Procedure

5

A 2 cm piece of dialysis tubing (molecular weight cut-off 12-14,000, 10 mm flat width) containing matrix metalloproteinase enzyme (stromelysin, collagenase or gelatinase in 0.5 mL of buffer) is implanted either ip or sc (in the back) of a rat (Sprague-Dawley, 150-200g) or mouse (CD-1, 25-50g) under anesthesia. Drugs are administered PO, IP, SC or IV through a canula in the jugular vein. Drugs are administered in a dose volume of 0.1 to 0.25 mL/animal. Contents of the dialysis tubing is collected and enzyme activity assayed.

Enzyme reaction rates for each dialysis tube are calculated. Tubes from at least 3 different animals are used to calculate the mean \pm sem. Statistical significance ($p < 0.05$) of vehicle-treated animals versus drug-treated animals is determined by analysis of variance. (*Agents and Actions* 21: 331, 1987).

Test Procedure for Measuring TACE Inhibition

Using 96-well black microtiter plates, each well receives a solution composed of 10 μ L TACE (final concentration 1 μ g/mL), 70 μ L Tris buffer, pH 7.4 containing 10% glycerol (final concentration 10 mM), and 10 μ L of test compound solution in DMSO (final concentration 1 μ M, DMSO concentration $< 1\%$) and incubated for 10 minutes at room temperature. The reaction is initiated by addition of a fluorescent peptidyl substrate (final concentration 100 μ M) to each well and then shaking on a shaker for 5 sec.

The reaction is read (excitation 340 nm, emission 420 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line. The slope of the line is calculated and represents the reaction rate.

The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($x \pm \text{sem}$) of the control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generate using multiple doses of drug and IC₅₀ values with 95% CI are estimated using linear regression.

- 68 -

Results of the above in-vitro and in-vivo matrix metalloproteinase inhibition and TACE inhibition standard pharmacological test procedures are given in Table I below.

5 Table I. Inhibition of MMP and TACE

	Example	MMP-1 ¹	MMP-9 ¹	MMP-13 ¹	TACE ¹	in-vivo MMP ²
	11	172	11	7	>1000	
	12	933	2	1	190	
	13	82	15	9	3%	
	14	108	8	6	24%	
10	15	139	25	12	7%	
	16	99	6	3	36%	64%(100)
	17	3100	8	16	401	
	18	152		26	627	
	19	194	2	4	314	
15	20	344	6	9	589	
	29	200	5	4		
	30	22	11	467	47	
	31	225	2	2	80	
	32	456	1	1	24	
20	33	1012	1	1		
	34	301	9	12	20	
	35	234	4	5	49	
	36	46	2	1	226	81%(50)
	37	65	2	1	124	
25	38	100	4	3	336	
	39	75	2	2	53	
	40	151	3	4	120	
	41	136	2	2	161	65%(50)
	42	5200	874	37	16%	
30	43	43%	71%	63%	20%	
	44	65%	59%	73%	5%	
	52	45	2.4	1.4	236	74%(100)
	60	39	2.9	2.5	160	53%(50)

Table I (continued)

Example	MMP-1 ¹	MMP-9 ¹	MMP-13 ¹	TACE ¹	in-vivo MMP ²
61	36	2.3	2.3	214	72%(50)
62	1236	5.7	23	46%	
63	721	6.8	23		
5 64	913	5.5	19		
66	512	0.81	0.27		5%(25)
67	96	3.0	2.4	138±11	
68	96	3.0	2.4	115±5	80%(50)
86	131	6.9	10.0	118±6	
10 87	111	8.9	10	147±5	12%(25)
88	643	10.4	18.7	40.5%	
89	116	0.8	1.1	356±15	6%(25)
90	550	2.3	3.0	535±77	39%(25)
91(HCl)	1805	1.8	1.1	38.7%	54%(25)
15 92	60.2% at 10 µM	12.0	6.1	>1644	30%(25)
93	8786	10.1	3.9	34.9%	49%(25)
94	2548	3.0	3.2	41.6%	12%(25)

1. IC₅₀ nM or % inhibition at 1 µM concentration

20 2. % inhibition (dose, mg/kg), p.o. vs MMP-13

25 Based on the results obtained in the standard pharmacological test procedures described above, the compounds of this invention were shown to be inhibitors of the enzymes MMP-1, MMP-9, MMP-13 and TNF-α converting enzyme (TACE) and are therefore useful in the treatment of disorders such as arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection.

30 The compounds of this invention are also useful in treating or inhibiting pathological changes mediated by matrix metalloproteinases such as atherosclerosis, atherosclerotic plaque formation, reduction of coronary thrombosis from atherosclerotic plaque rupture, restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, angiogenesis, tumor metastasis, tumor growth,

- 70 -

osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, age
5 related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft rejection.

10

Compounds of this invention may be administered neat or with a pharmaceutical carrier to a patient in need thereof. The pharmaceutical carrier may be solid or liquid.

Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants,
15 compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active
20 ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers may be used in preparing solutions, suspensions, emulsions,
25 syrups and elixirs. The active ingredient of this invention can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents,
30 colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferable sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, and oils (e.g., fractionated coconut oil and arachis
35 oil). For parenteral administration the carrier can also be an oily ester such as ethyl

- 71 -

oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection.

5 Sterile solutions can also be administered intravenously. Oral administration may be either liquid or solid composition form.

The compounds of this invention may be administered rectally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of this invention may be formulated into an aqueous or
10 partially aqueous solution, which can then be utilized in the form of an aerosol. The compounds of this invention may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms
15 such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semi-solid emulsions of either the oil in water or water in oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such
20 as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

The dosage to be used in the treatment of a specific patient suffering a MMP or TACE dependent condition must be subjectively determined by the attending physician.
25 The variables involved include the severity of the dysfunction, and the size, age, and response pattern of the patient. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached. Precise dosages for oral, parenteral, nasal, or intrabronchial administration will be determined by the
30 administering physician based on experience with the individual subject treated and standard medical principles. Projected oral daily dosages are 2 - 500 mg/kg, preferred oral daily dosages are 2 - 50 mg/kg, and more preferred oral daily dosages are 5 - 25 mg/kg.

35 Preferably the pharmaceutical composition is in unit dosage form, e.g., as tablets or capsules. In such form, the composition is sub-divided in unit dose containing

- 72 -

appropriate quantities of the active ingredient; the unit dosage form can be packaged compositions, for example packed powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

- 73 -


What is claimed:

1. A compound having the formula:

5


B

wherein **B** is




10


P and Q are




or



P is




, Q is



, and vice versa;

- 15 T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R^1 ;
 Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided that no more than 2 of T, U, W, and X are nitrogen;

- 74 -




is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms

- which may contain 0-2 heteratoms selected from nitrogen, oxygen, and sulfur, in addition to any heteroatoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R^1 ;
- 5 Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteroatoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri- substituted with R^1 ;
- 10 R^1 is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, $-(CH_2)_n$, $-OR^2$, $-CN$, $-COR^2$, perfluoroalkyl of 1-4 carbon atoms, $-CONR^2R^3$, $-S(O)_xR^2$, $-OPO(OR^2)OR^3$, $-PO(OR^2)R^3$, $-OC(O)NR^2R^3$, $-COOR^2$, $-CONR^2R^3$, $-SO_3H$, $-NR^2R^3$, $-NR^2COR^3$, $-NR^2COOR^3$, $-SO_2NR^2R^3$, $-NO_2$, $-N(R^2)SO_2R^3$, $-NR^2CONR^2R^3$, $-NR^2C(=NR^3)NR^2R^3$, $-SO_2NHCOR^4$, $-CONHSO_2R^4$, $-tetrazol-$
- 15 $-5-yl$, $-SO_2NHCN$, $-SO_2NHCONR^2R^3$, or Z;
- V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteroatoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R^2 ;
- 20 R^2 and R^3 are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;
- R^4 is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;
- 25 R^5 is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V;
- $n = 1-6$;
- $x = 0-2$

- 75 -

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein



5

3. The compound according to claim 2, wherein


W and X are carbon; and

T is nitrogen;


U is carbon, optionally substituted with R¹

10 or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 3, wherein



15



substituted with R¹;

or a pharmaceutically acceptable salt thereof.

20

- 76 -

5. The compound according to claim 1 which is selected from the group consisting of:
- 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
 - 5 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-8-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-bromo-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-7-bromo-quinoline-3-carboxylic acid hydroxyamide,
 - 10 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[(4-methoxybenzenesulfonyl)-pyridin-3-ylmethylamino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
 - 15 4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-t-butyl-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-methyl-quinoline-3-carboxylic acid hydroxyamide,
 - 8-Ethyl-4-[benzyl-(4-methoxybenzenesulfonyl)-amino]-quinoline-3-carboxylic acid hydroxyamide,
 - 20 4-[Benzyl-(4-methoxybenzenesulfonyl)-amino]-8-(1-methylethyl)-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-8-vinyl-quinoline-3-carboxylic acid hydroxyamide,
 - 25 4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-6-nitro-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]-8-bromo-quinoline-3-carboxylic acid hydroxyamide,
 - 4-[Methyl-[4-(pyridin-4-yloxy)-benzenesulfonyl]-amino]-6-iodo-quinoline-3-carboxylic acid hydroxyamide,
 - 30 4-[Methyl-(4-(pyridin-4-yloxy)-benzenesulfonyl)-amino]-6-iodo-quinoline-3-carboxylic acid hydroxyamide hydrochloride,
 - 4-[Ethyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethynyl-quinoline-3-carboxylic acid hydroxyamide,

- 4-[Methyl-(4-methoxy-benzenesulfonyl)-amino]-6-phenylethyl-quinoline-3-carboxylic acid hydroxyamide,
- 4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-methoxy-quinoline-3-carboxylic acid hydroxyamide,
- 5 4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-bromo-quinoline-3-carboxylic acid hydroxyamide,
- 4-[(4-methoxy-benzenesulfonyl)-pyridin-3-ylmethyl amino]-8-Benzyl-quinoline-3-carboxylic acid hydroxyamide,
- 4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-iodo-quinoline-3-carboxylic acid hydroxyamide,
- 10 4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-phenyl-quinoline-3-carboxylic acid hydroxyamide,
- 4-[(4-Methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-8-thiophen-2-yl-quinoline-3-carboxylic acid hydroxyamide,
- 15 4-[(Biphenyl-4-sulfonyl)-pyridin-3-ylmethyl-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
- 4-[(Octane-1-sulfonyl)-pyridin-3-ylmethyl-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
- 4-[Pyridin-3-ylmethyl-(toluene-4-sulfonyl)-amino]-7-trifluoromethyl-quinoline-3-carboxylic acid hydroxyamide,
- 20 4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
- 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
- 25 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide hydrochloride,
- 4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
- 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
- 30 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide, hydrochloride,
- 4-[Benzyl-(4-methoxybenzenesulfonyl)amino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,

- 78 -

- 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-phenyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide, hydrochloride,
5 4-[(4-Methoxybenzenesulfonyl)pyridin-2-ylmethyl amino] -1,3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)pyridin-4-ylmethylamino] -1,3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1-isopropyl - 1H-pyrazolo[3,4b]pyridin -5-carboxylic acid, hydroxyamide,
10 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1-benzyl-1H-pyrazolo[3,4-b]pyridin -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)amino] -1-benzyl -3-methyl -1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
15 4-[(4-Methoxybenzenesulfonyl) 2-thienylmethylamino] -1, 3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl) -3-thienylmethylamino] -1, 3- dimethyl - 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl) pyridin-3-ylmethylamino] -1- (2,4- dimethoxyphenyl) -
20 3-methyl- 1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino] -1-(2- methoxyphenyl)-3-methyl-1H-pyrazolo[3,4-b]pyridine -5-carboxylic acid, hydroxyamide,
4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino} -1,3-dimethyl-1H-pyrazolo[3,4-b] pyridine-5-carboxylic acid, hydroxyamide,
25 4-{Methyl-[4-(phenoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
4-[Methyl-(4-methoxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
4-[Methyl-(4-propyloxybenzenesulfonyl)amino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
30 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-ethyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,

- 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-tert-butyl-3-methyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-1-methyl-3-tert-butyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 5 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 4-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 7-[(4-Methoxybenzenesulfonyl)pyridin-3-ylmethylamino]-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid, hydroxyamide,
 10 4-[(4-(4-Chlorophenoxy)benzenesulfonyl)methylamino]-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 4-[(4-(4-Chlorophenoxy)benzenesulfonyl)methylamino]-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 15 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-3-methylisothiazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-1-methyl-3-phenyl-1H-pyrazolo[3,4-b]pyridine-5-carboxylic acid, hydroxyamide, and
 20 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-3-methylisoxazolo[5,4-b]pyridine-5-carboxylic acid, hydroxyamide,
 or a pharmaceutically acceptable salt thereof.
- 25 6. The compound according to claim 1 which is selected from the group consisting of:
 7-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-2-methylpyrazolo[1,5-a]pyrimidine-6-carboxylic acid, hydroxyamide,
 4-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-7-methyl-1,8-naphthyridine-3-carboxylic acid, hydroxyamide,
 30 7-{Methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-2,3-dimethylimidazo[4,5-b]pyridine-6-carboxylic acid, hydroxyamide,
 2-Methyl-4-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}thieno[3,4-b]pyridine-3-carboxylic acid, hydroxyamide, and

- 80 -

5-Methyl-7-{methyl-[4-(4-pyridinyloxy)benzenesulfonyl]amino}-thieno[3.2-b]pyridine-6-carboxylic acid, hydroxyamide, or a pharmaceutically acceptable salt thereof.


5

7. A method of inhibiting pathological changes mediated by matrix metalloproteinases in a mammal in need thereof which comprises administering to said mammal a therapeutically effective amount of a compound having the formula:



10

B

wherein **B** is




15



20 T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R¹;

Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided that no more than 2 of T, U, W, and X are nitrogen;

5



is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms

which may contain 0-2 heteroatoms selected from nitrogen, oxygen, and sulfur, in addition to any heteroatoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R¹;

10 Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteroatoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri- substituted with R¹;

15 R¹ is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -(CH₂)_nZ, -OR², -CN, -COR², perfluoroalkyl of 1-4 carbon atoms, -CONR²R³, -S(O)_xR², -OPO(OR²)OR³, -PO(OR²)R³, -OC(O)NR²R³, -COOR², -CONR²R³, -SO₃H, -NR²R³, -NR²COR³, -NR²COOR³, -SO₂NR²R³, -NO₂, -N(R²)SO₂R³, -NR²CONR²R³, -NR²C(=NR³)NR²R³, -SO₂NHCOR⁴, -CONHSO₂R⁴, -tetrazol-
20 5-yl, -SO₂NHCN, -SO₂NHCONR²R³, or Z;

V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteroatoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R²;

25 R² and R³ are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

R⁴ is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

- 82 -

R⁵ is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V;

n = 1-6;

x = 0-2

5 or a pharmaceutically acceptable salt thereof.

8. The method according to claim 7 wherein the matrix metalloproteinase mediated condition treated is atherosclerosis, atherosclerotic plaque formation, reduction of coronary thrombosis from atherosclerotic plaque rupture, restenosis, MMP-mediated
10 osteopenias, inflammatory diseases of the central nervous system, skin aging, angiogenesis, tumor metastasis, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease
15 of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, or periodontal disease.

9. The method according to claim 7, wherein the matrix metalloproteinase mediated condition treated is age related macular degeneration, diabetic retinopathy, proliferative
20 vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft rejection.




10. A method of inhibiting pathological changes mediated by TNF- α converting
25 enzyme (TACE) in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound having the formula:

B

30

wherein **B** is

- 83 -




5

T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R¹;

Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided that no more than 2 of T, U, W, and X are

10



is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms

- 84 -

which may contain 0-2 heteroatoms selected from nitrogen, oxygen, and sulfur, in addition to any heteroatoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R¹;

5 Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteroatoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri- substituted with R¹;

10 R¹ is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, $-(CH_2)_nZ$, $-OR^2$, $-CN$, $-COR^2$, perfluoroalkyl of 1-4 carbon atoms, $-CONR^2R^3$, $-S(O)_xR^2$, $-OPO(OR^2)OR^3$, $-PO(OR^2)R^3$, $-OC(O)NR^2R^3$, $-COOR^2$, $-CONR^2R^3$, $-SO_3H$, $-NR^2R^3$, $-NR^2COR^3$, $-NR^2COOR^3$, $-SO_2NR^2R^3$, $-NO_2$, $-N(R^2)SO_2R^3$, $-NR^2CONR^2R^3$, $-NR^2C(=NR^3)NR^2R^3$, $-SO_2NHCOR^4$, $-CONHSO_2R^4$, $-tetrazol-5-yl$, $-SO_2NHCN$, $-SO_2NHCONR^2R^3$, or Z;

15 V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteroatoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R²;

20 R² and R³ are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

R⁴ is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

25 R⁵ is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V;

n = 1-6;

x = 0-2

or a pharmaceutically acceptable salt thereof.

30 11. The method according to claim 11 wherein the condition treated is rheumatoid arthritis, graft rejection, cachexia, anorexia, inflammation, fever, insulin resistance, septic shock, congestive heart failure, inflammatory disease of the central nervous system, inflammatory bowel disease, or HIV infection.




- 85 -

15. A pharmaceutical composition comprising a compound having the formula:

B

5

wherein **B** is




10

T, U, W, and X are each, independently, carbon or nitrogen, provided that when T or U is carbon, either may be optionally substituted with R¹;

15 Y is carbon, nitrogen, oxygen or sulfur, provided that at least one of T, U, W, X, and Y is not carbon, and further provided that no more than 2 of T, U, W, and X are nitrogen;

- 86 -



is a phenyl ring or is a heteroaryl ring of ring 5-6 atoms

which may contain 0-2 heteratoms selected from nitrogen, oxygen, and sulfur, in addition to any heteratoms defined by W or X; wherein the phenyl or heteroaryl ring may be optionally mono-, di-, or tri- substituted with R¹;

5 Z is a phenyl, naphthyl, heteroaryl, or heteroaryl fused to phenyl, wherein the heteroaryl moiety contains of 5-6 ring atoms and 1-3 heteratoms selected from nitrogen, oxygen, or sulfur; wherein the phenyl, naphthyl, heteroaryl, or phenyl fused heteroaryl moieties may be optionally mono-, di-, or tri- substituted with R¹;

10 R¹ is hydrogen, halogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -(CH₂)_nZ, -OR², -CN, -COR², perfluoroalkyl of 1-4 carbon atoms, -CONR²R³, -S(O)_xR², -OPO(OR²)OR³, -PO(OR²)R³, -OC(O)NR²R³, -COOR², -CONR²R³, -SO₃H, -NR²R³, -NR²COR³, -NR²COOR³, -SO₂NR²R³, -NO₂, -N(R²)SO₂R³, -NR²CONR²R³, -NR²C(=NR³)NR²R³, -SO₂NHCOR⁴, -CONHSO₂R⁴, -tetrazol-5-yl, -SO₂NHCN, -SO₂NHCONR²R³, or Z;

V is a saturated or partially unsaturated heterocycloalkyl ring of 5-7 ring atoms having 1-3 heteratoms selected from N, O, or S, which may be optionally mono-, or di-substituted with R²;

20 R² and R³ are each, independently, hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

R⁴ is alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms, Z or V;

25 R⁵ is hydrogen, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, Z, or V;

n = 1-6;

x = 0-2

or a pharmaceutically acceptable salt thereof, and a pharmaceutical carrier.

INTERNATIONAL SEARCH REPORT

Intr onal Application No
PCT/US 98/07380

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D215/54 A61K31/63 A61K31/47 A61K31/44 C07D401/12
C07D471/04 C07D513/04 C07D498/04 C07D495/04 C07F9/547
A61K31/66 //(C07D513/04,277:00,221:00),(C07D498/04,261:00,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 18194 A (HOECHST) 22 May 1997 see page 17, paragraph 4 - page 18, paragraph 1; claim 1	1, 15
P, A	EP 0 803 505 A (ADIR) 29 October 1997 see page 3, line 3 - line 17	1, 15
E	WO 98 16514 A (AMERICAN CYANAMID) 23 April 1998 see claims 1, 10	1, 15

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 July 1998

Date of mailing of the international search report

03/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

Inte onal Application No
PCT/US 98/07380

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 221:00), (C07D471/04, 231:00, 221:00), (C07D513/04, 275:00, 221:00), (C07D471/04, 221:00, 221:00), (C07D471/04, 235:00, 221:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "8" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
24 July 1998		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx 31 651 epo nl. Fax: (+31-70) 340-3018		Authorized officer
		Alfaro Faus, I

Form PCT:ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/07380

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7 to 11
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 7 to 11
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/07380

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9718194 A	22-05-1997	DE 19542189 A DE 19612298 A AU 7562496 A	15-05-1997 02-10-1997 05-06-1997
EP 803505 A	29-10-1997	FR 2748026 A AU 1912197 A CA 2203618 A CN 1165817 A JP 10059936 A NO 971862 A PL 319684 A	31-10-1997 30-10-1997 26-10-1997 26-11-1997 03-03-1998 27-10-1997 27-10-1997
WO 9816514 A	23-04-1998	AU 4980697 A	11-05-1998

Form PCT/ISA/210 (patent family annex) (July 1992)